

Manuscript Number: 20161231R1

Title: Handgrip strength and ideal cardiovascular health among Colombian children and adolescents: The FUPRECOL Study

Article Type: Original Article

Keywords: Muscular strength; Risk factors; Cardiovascular health; Health behaviors; Prevention.

Corresponding Author: Dr. Robinson Ramírez-Vélez, Ph.D

Corresponding Author's Institution: Universidad del Rosario

First Author: ROBINSON RAMIREZ-VELEZ, Ph.D

Order of Authors: ROBINSON RAMIREZ-VELEZ, Ph.D; Alejandra Tordecilla-Sanders, MSc; Jorge E Correa-Bautista, Ph.D; Mark D Peterson, Ph.D; Antonio Garcia-Hermoso, Ph.D

Abstract: Objective: To evaluate the association between handgrip strength and ideal cardiovascular health in Colombian children and adolescents.

Study design: During the 2014-2015 school years, we examined a cross-sectional component of the FUPRECOL study. Participants included 1,199 (n=627 boys) youths from Bogota (Colombia). Handgrip strength was measured with a standard adjustable hand held dynamometer and expressed relative to body mass (HG/body mass) and as absolute values in kilograms. Ideal cardiovascular health, as defined by the American Heart Association, was determined as meeting ideal levels of the following components: four behaviours (smoking status, body mass index, cardiorespiratory fitness, and diet) and three factors (total cholesterol, blood pressure and glucose).

Results: Higher levels of handgrip strength (both absolute and relative values) were associated with a higher frequency of ideal cardiovascular health metrics in both sexes (p for trend  $\leq 0.001$ ). Also, higher levels of handgrip strength were associated with a greater number of ideal health behaviours (p for trend  $< 0.001$  in both boys and girls), and with a higher number of ideal health factors in boys (p for trend  $< 0.001$ ). Finally, levels of handgrip strength were similar between ideal versus non-ideal glucose or total cholesterol groups in girls.

Conclusions: Handgrip strength was strongly associated with ideal cardiovascular health in Colombian children and adolescents, and thus supports the relevance of early targeted interventions to promote strength adaptation and preservation as part of primordial prevention.

1 **Title:** Handgrip strength and ideal cardiovascular health among Colombian children and  
2 adolescents: The FUPRECOL Study

3 **Running title:** Handgrip strength and cardiovascular health in Colombian youth

4 Robinson Ramírez-Vélez, PhD<sup>1</sup>, (E:mail: [robinson.ramirez@urosario.edu.co](mailto:robinson.ramirez@urosario.edu.co))

5 Alejandra Tordecilla-Sanders, Msc<sup>1</sup>, (E:mail: [alesanders\\_0615@hotmail.com](mailto:alesanders_0615@hotmail.com))

6 Jorge Enrique Correa-Bautista, PhD<sup>1</sup>, (E:mail: [jorge.correa@urosario.edu.co](mailto:jorge.correa@urosario.edu.co))

7 Mark D. Peterson, PhD<sup>2,3</sup>, (E:mail: [mdpeterz@med.umich.edu](mailto:mdpeterz@med.umich.edu))

8 Antonio Garcia-Hermoso, PhD<sup>4,5</sup> (E:mail: [antonio.garcia.h@usach.cl](mailto:antonio.garcia.h@usach.cl))

9

10 <sup>1</sup> Centro de Estudios para la Medición de la Actividad Física «CEMA». Escuela de  
11 Medicina y Ciencias de la Salud, Universidad del Rosario, Bogotá D.C, Colombia.

12 <sup>2</sup> Department of Physical Medicine and Rehabilitation, University of Michigan, Ann Arbor,  
13 Michigan, United States.

14 <sup>3</sup> Global REACH, University of Michigan Medical School, University of Michigan, Ann  
15 Arbor, Michigan, United States.

16 <sup>4</sup> Laboratorio de Ciencias de la Actividad Física, el Deporte y la Salud, Universidad de  
17 Santiago de Chile, USACH, Santiago, Chile.

18 <sup>5</sup> Facultad de Ciencias de la Actividad Física, Universidad San Sebastián, Santiago, Chile.

19

20 **Reprint requests:** Robinson Ramírez-Vélez, PhD. Universidad del Rosario, Bogotá, D.C,  
21 Colombia. Phone: +57 (1) 2970200 ext. 3428, E-mail: [robin640@hotmail.com](mailto:robin640@hotmail.com) //  
22 [robinson.ramirez@urosario.edu.co](mailto:robinson.ramirez@urosario.edu.co)

23

24 **Robinson Ramírez-Vélez, PhD:** Wrote the first draft of the manuscript.

25

26 **Statement of honorarium, grant, or other form of payment:** The FUPRECOL Study  
27 was carried out with the financial support of Instituto Colombiano para el Desarrollo de la  
28 Ciencia y la Tecnología “Francisco José de Caldas” COLCIENCIAS (Contract N° 671-  
29 2014 Code 122265743978). This article presents independent research commissioned by  
30 COLCIENCIAS under its Programme Grants for Applied Research funding scheme  
31 (Convocatoria 671-2014). The content of this paper reflects the author’s views alone, and  
32 the Colombian Community or the COLCIENCIAS is not liable for any use that may be  
33 made of the information contained herein.

34

35 **Key words:** Muscular strength; Risk factors; Cardiovascular health; Health behaviors;  
36 Prevention.

37 **Competing interests:** The authors declare that they have no competing interests.

1 **ABSTRACT**

2 **Objective:** To evaluate the association between handgrip strength and ideal  
3 cardiovascular health in Colombian children and adolescents.

4 **Study design:** During the 2014–2015 school years, we examined a cross-sectional  
5 component of the FUPRECOL study. Participants included 1,199 (n=627 boys) youths  
6 from Bogota (Colombia). Handgrip strength was measured with a standard adjustable hand  
7 held dynamometer and expressed relative to body mass (HG/body mass) and as absolute  
8 values in kilograms. Ideal cardiovascular health, as defined by the American Heart  
9 Association, was determined as meeting ideal levels of the following components: four  
10 behaviours (smoking status, body mass index, cardiorespiratory fitness, and diet) and three  
11 factors (total cholesterol, blood pressure and glucose).

12 **Results:** Higher levels of handgrip strength (both absolute and relative values) were  
13 associated with a higher frequency of ideal cardiovascular health metrics in both sexes (p  
14 for trend  $\leq 0.001$ ). Also, higher levels of handgrip strength were associated with a greater  
15 number of ideal health behaviours (p for trend  $< 0.001$  in both boys and girls), and with a  
16 higher number of ideal health factors in boys (p for trend  $< 0.001$ ). Finally, levels of  
17 handgrip strength were similar between ideal versus non-ideal glucose or total cholesterol  
18 groups in girls.

19 **Conclusions:** Handgrip strength was strongly associated with ideal cardiovascular  
20 health in Colombian children and adolescents, and thus supports the relevance of early  
21 targeted interventions to promote strength adaptation and preservation as part of primordial  
22 prevention.

## 1 INTRODUCTION

2 Low muscular strength, as determined by handgrip dynamometry, is a recognized  
3 marker of poor health during adolescence<sup>1,2</sup>, and is associated with disease and mortality in  
4 adulthood<sup>3-5</sup>. Numerous studies support an inverse relationship between muscular strength  
5 and cardiovascular disease (CVD) risk factors in young populations, and generally express  
6 muscular strength relative to body mass<sup>1,6</sup>. Epidemiological studies indicate that muscle  
7 weakness has been associated with a higher frequency of adverse health consequences  
8 including obesity, systemic low-grade inflammation, and insulin resistance<sup>7-10</sup>. CVD events  
9 occur most frequently during or after the fifth decade of life, and yet the precursors of  
10 disease originate in childhood and adolescence<sup>1,11</sup>.

11 In response to the increasing burden of CVD risk factors, the American Heart  
12 Association established several strategic goals<sup>12</sup>. In 2010, the American Heart Association  
13 released a set of cardiovascular health metrics for adults and children that were intended to  
14 prioritize cardiovascular health, as opposed to cardiovascular disease<sup>12</sup>. Population-  
15 representative studies have shown a low prevalence of ideal cardiovascular health (CVH)  
16 metrics in U.S. children and adolescents, particularly for achieving physical activity  
17 recommendations and dietary intake<sup>13,14</sup>. Data from the Cardiovascular Risk in Young  
18 Finns Study and The Healthy Lifestyle in Europe by Nutrition also demonstrated that  
19 children and adolescence with a higher number of ideal CVH components had a reduced  
20 risk for hypercholesterolemia, hypertension, and elevated blood glucose<sup>15</sup>. Increases in  
21 ideal CVH are directly associated with aortic elasticity<sup>16</sup> and healthier levels of  
22 cardiorespiratory fitness in adolescents<sup>17</sup>. Among adults, a recent systematic review<sup>18</sup>  
23 reported an inverse association between number of ideal CVH metrics and early all-cause

1 and CVD-related mortality. Improved understanding of the health-risks associated with  
2 muscle weakness will help to inform the development of targeted interventions for different  
3 phenotypes.

4 Obesity and physical inactivity are leading CVD risk factors among Hispanic/Latino  
5 adults, raising concerns about whether an increased risk of these conditions also is  
6 manifested at younger ages<sup>19</sup>. Previous research has demonstrated an independent  
7 association between muscle weakness and increased cardiometabolic risk factors<sup>20,21</sup>.

8 In Colombia, a region which has undergone a well-documented epidemiologic  
9 transition and epidemic of CVD<sup>19-22</sup>, relatively little research on physical activity<sup>22</sup> and  
10 physical fitness exists<sup>23,24</sup>. Therefore, describing the magnitude of these risk factors in  
11 youth is important for prioritizing prevention and public health efforts<sup>19</sup>. Nevertheless,  
12 there have been no studies to date to determine the association between handgrip strength  
13 and ideal CVH in Latin American youth. Therefore, the objective of the present study was  
14 to investigate the relationship between handgrip strength and ideal CVH among Colombian  
15 children and adolescents.

16

## 17 **METHODS**

### 18 *Participants and Study Design*

19 This study aimed to examine the relationships between physical fitness levels,  
20 healthy and unhealthy behaviors, and cardiometabolic risk factors in Colombian children  
21 and adolescents. During the 2014–2015 school years, we examined a cross-sectional  
22 component of the FUPRECOL study<sup>25-27</sup> (*in Spanish*, ASOCIACIÓN DE LA **FUERZA**  
23 **PRENSIL** CON MANIFESTACIONES DE RIESGO CARDIOVASCULAR  
24 TEMPRANAS EN NIÑOS Y ADOLESCENTES **COLOMBIANOS**). The sample

1 consisted of children and adolescents (boys  $n = 4,000$  and girls  $n = 4,000$ ) aged 9–17.9  
2 years. Blood sampling was randomly performed in one-third of the recruited subjects  
3 ( $n=2,775$ ). From this subgroup, 1,199 schoolchildren (52.2% boys) had valid data muscular  
4 strength and all components included in the ideal CVH concept. There were no differences  
5 in the study key characteristics (i.e., age, sex distribution, BMI, and muscular strength)  
6 between the current study sample and the original FUPRECOL Study sample ( $n=8,000$ , all  
7  $p>0.100$ ). The children and adolescents were of low to middle socioeconomic status (SES,  
8 1–3 defined by the Colombian government), enrolled in public elementary and high schools  
9 (grades 5 through 11), and from the capital district of Bogota in a municipality in the  
10 Cundinamarca Department in the Andean region. A convenience sample of volunteers was  
11 included and grouped by sex and age with 1-year increments (a total of 9 groups).

## 12 ***Measurements***

### 13 *Handgrip strength assessment*

14 Consistent with recommendations<sup>28,29</sup>, we restricted our analysis to the following  
15 health-related<sup>30</sup> field-based tests that have demonstrated adequate levels of criterion-related  
16 validity, and reliability<sup>27-29</sup> in the assessment of two dimensions of muscular strength:  
17 handgrip strength and normalized handgrip strength in kg/body mass in kg<sup>31-33</sup>. Handgrip  
18 was measured using a standard adjustable hand held dynamometer (Takei Digital Grip  
19 Strength Dynamometer Model T.K.K.540<sup>®</sup>, Takei Scientific Instruments Co., Ltd, Niigata,  
20 Japan). Participants were given a brief demonstration and verbal instructions for the test,  
21 and if necessary, the dynamometer was adjusted to the participant's hand size according to  
22 predetermined protocols<sup>27</sup>. Handgrip strength was measured with the subject in a standing  
23 position, with the shoulder adducted and neutrally rotated, and arms parallel but not in  
24 contact with the body. The participants were asked to squeeze the handle as hard as

1 possible for a maximum of 3-5 seconds, and no verbal encouragement was given during the  
2 test. Handgrip strength performance was recorded as the best score from either hand,  
3 without consideration for hand dominance. Since there is substantial covariance between  
4 strength capacity and body mass—and, moreover, the links between muscle strength and  
5 both physical function and chronic health are mediated by the proportion of strength  
6 relative to body mass—grip strength was normalized as strength per body mass [i.e. (grip  
7 strength in kg)/(body mass in kg)]. Handgrip measurements in a subsample (n=229, median  
8 age = 12.8 ±2.4 y, 46.2±12.4 kg, 1.50±0.1 m, 19.9±3.1 kg/m<sup>2</sup>) were recorded to ensure  
9 reproducibility on the day of the study. The reproducibility of our data was R=0.96. Intra-  
10 rater reliability was assessed by determining the intraclass correlation coefficient (0.98, CI  
11 95% 0.97 to 0.99). Monthly, each dynamometer was tested using a standardized calibration  
12 procedure which showed that the device was within 1 kg of accuracy over the whole  
13 measuring range (from 0 to 100 kg), and with a 100 g sensitivity.

#### 14 *Anthropometric measurements*

15         Body weight was measured in the subjects' underwear and with no shoes, using  
16 electronic scales (Tanita<sup>®</sup> BC544, Tokyo, Japan) with a low technical error of measurement  
17 (Technical error of measurement = 0.510%). Height was measured using a mechanical  
18 stadiometer platform (Seca<sup>®</sup> 274, Hamburg, Germany; Technical error of measurement =  
19 0.01%). Body mass index (BMI) was calculated as the body weight in kilograms divided by  
20 the square of height in meters (kg/m<sup>2</sup>). Obesity status was defined as having a BMI above  
21 the age and sex-specific thresholds of the International Obesity Task Force (IOTF)<sup>34</sup>.  
22 Participants who had a BMI <85<sup>th</sup> percentile were categorized as meeting the ideal  
23 cardiovascular health criteria for BMI.

#### 24 *Biochemical determinations*

1 Blood samples were collected between 6:00 and 8:00 am by two experienced  
2 paediatric phlebotomists after at least 12 hours fasting. Before the extraction, fasting  
3 condition was confirmed by the child and parents. Blood samples were obtained from an  
4 antecubital vein, and analyses were subsequently completed within 1 day from collection.  
5 In children and adolescence, levels of total cholesterol have been defined as “ideal” <4.40  
6 mmol/L (<170 mg/dL), or “non-ideal”  $\geq 4.40$  mmol/L ( $\geq 170$  mg/dL). Fasting serum glucose  
7 concentrations were analysed enzymatically and also classified as ideal <5.6 mmol/L (<100  
8 mg/ dL), or non-ideal  $\geq 5.6$  mmol/L ( $\geq 100$  mg/dL). Inter-assay reproducibility (coefficient  
9 of variation) was determined from 80 replicate analyses of 8 plasma pools over 15 days,  
10 and shown to be 2.6% for total cholesterol and 1.5% for serum glucose. None of the study  
11 youths were on any drug treatments.

#### 12 *Resting blood pressure*

13 Blood pressure was measured using an electronic oscillometric device, (Riester Ri-  
14 Champion model, Jungingen, Germany) after being seated in a quiet room for 10 min with  
15 their back supported and feet on the ground. Two blood pressure readings were taken with a  
16 10 min interval of quiet rest. Before blood pressure session monitoring, the accuracy of the  
17 device was tested against a standard mercury sphygmomanometer in a random sub-sample  
18 (n=25) to ensure that there was no consistent difference of >10 mm Hg in measured blood  
19 pressure; and inter-observer variability was R=0.96. Mean systolic blood pressure was  
20 defined as ideal (<90<sup>th</sup> centile *and* mean diastolic blood pressure <90<sup>th</sup> centile), or non-ideal  
21 (systolic blood pressure  $\geq 90^{\text{th}}$  centile *or* diastolic blood pressure  $\geq 90^{\text{th}}$  centile). All centile-  
22 based threshold limits were sex- and age-specific and selected on the basis of the  
23 International Diabetes Federation<sup>35</sup> and the modified De Ferranti et al.<sup>36</sup> definitions of  
24 metabolic syndrome.



## 1 *Dietary Assessment*

2           Dietary intake and food consumption was assessed by the Kidmed questionnaire<sup>37</sup>.  
3 This tool consists of sixteen questions related to the principles of Mediterranean dietary  
4 patterns. The score ranges from -4 to 12 points, since questions with negative connotations  
5 with respect to the Mediterranean diet are assigned a value of -1 (frequent intake of fast  
6 food, increased consumption of sweets, skipping breakfast, frequent intake of pastries for  
7 breakfast). Parameters with positive connotations are assigned + 1 point (e.g. takes a fruit  
8 or fruit juice every day, consumes fish regularly (at least 2-3 times/week)) as indicated  
9 previously<sup>37</sup>. As suggested by Serra-Majem et al.<sup>37</sup>, the total score was divided into three  
10 categories of Mediterranean diet quality: (1)  $\leq 3$  points = poor diet quality; (2) 4-7 points =  
11 average diet quality; and (3)  $\geq 8$  points = good diet quality (optimal Mediterranean diet  
12 style). Participants who had at least  $\geq 8$  points were categorised as having an ideal healthy  
13 diet, whereas children and adolescents with  $< 7$  points were classified as having a non-ideal  
14 healthy diet.

## 15 *Cardiorespiratory fitness*

16           Although the American Heart Association relied on physical activity to determine  
17 active habits, we used estimated cardiorespiratory fitness (CRF) due to its robust  
18 association with cardiovascular risk factors<sup>38</sup>, and ideal CVH<sup>12,17</sup> in this population. We  
19 estimated CRF with the 20 m shuttle run test, as previously described by Leger et al.<sup>39</sup>  
20 Participants ran in a straight line between two lines 20 m apart, while keeping pace with  
21 pre-recorded audio signals. The initial speed was 8.5 km/hr and increased by 0.5 km/hr  
22 each minute. The test was finished when the participant failed to reach the end lines,  
23 keeping pace with the audio signals on two consecutive occasions or when the subject  
24 stopped because of fatigue. Results were recorded to the nearest stage (minute) completed.

1 Healthy cardiorespiratory fitness was defined by using either the cut-off by sex- and age  
2 (shuttle-runs or estimated  $VO_{2peak}$ ) listed in the healthy fitness zone (needs improvement  
3 and health risk). The FITNESSGRAM<sup>®40</sup> has been shown to have cardio-metabolic health  
4 predictive value<sup>41</sup>, and  $VO_{2peak}$  cut-off points were validated against the presence of  
5 metabolic syndrome using nationally representative U.S. data<sup>41,42</sup>.

### 6 *Smoking habits*

7 Data on smoking were collected via self-reported questionnaires [number of  
8 cigarettes smoked per day]. Students who reported that they had never smoked were  
9 categorised as having an ideal smoking status and those who reported having smoked one  
10 or more cigarettes were categorized as presenting a non-ideal smoking status.

### 11 *Sexual maturation*

12 Sexual maturation was classified based on Tanner staging<sup>43</sup>, which uses self-  
13 reported puberty status to classify participants into stages I to V<sup>44</sup>. Each volunteer entered  
14 an isolated room where they categorized the development of their own genitalia (for boys),  
15 breasts (for girls), armpits (for boys) and pubic hair (for both genders) using a set of images  
16 exemplifying the various stages of sexual maturation. The reproducibility of our data  
17 reached  $R=0.78$ .

### 18 *Ideal cardiovascular health*

19 The metrics for ideal cardiovascular health in children and adolescents defined by  
20 the American Heart Association<sup>12</sup> were followed as precisely as possible (Table I, available  
21 at [www.jpeds.com](http://www.jpeds.com)). Finally, each participants was categorised into 5 health levels based  
22 on the number of ideal CVH metrics in the ideal range: the healthiest level (favourable  
23 ideal CVH score) was defined as having  $\geq 5$  metrics, the intermediate levels as 2 to 4  
24 metrics in the ideal range, and the unfavourable level as having 0-1 ideal CVH metrics. We

1 collapsed 0 with 1 and 5 with 7 ideal metrics due to relatively few youths who had 0 (2% of  
2 total cohort) or 6 (8% of total cohort) and 7 (2% of total cohort) ideal CVH metrics.

### 3 *Ethics Statement*

4 The Review Committee for Research on Human Subjects at the University of Rosario  
5 [Code N° CEI-ABN026-000262] approved all of the study procedures. A comprehensive  
6 verbal description of the nature and purpose of the study and its experimental risks was  
7 given to the participants and their parents/guardians. Written informed consent was  
8 obtained from parents and subjects before participation in the study. The protocol was in  
9 accordance with the latest revision of the Declaration of Helsinki and current Colombian  
10 laws governing clinical research on human subjects (Resolution 008430/1993 Ministry of  
11 health).

### 12 *Statistical analysis*

13 The data are presented as means, standard deviations, and percentages. The t-test  
14 was used to compare unadjusted means by sex. Differences on handgrip strength (both  
15 absolute and normalized handgrip strength in kg/body mass in kg) between ideal and non-  
16 ideal cardiovascular health components were assessed by analysis of covariance  
17 (ANCOVA), with handgrip strength as a dependent variable, the cardiovascular health  
18 component (ideal vs non-ideal) entered as a fixed factor, age as a covariate, and sexual  
19 maturation as a random factor. The association between handgrip strength and ideal CVH  
20 metrics, as well as with ideal CVH behaviours and factors separately, was assessed by  
21 ANCOVA as explained above. Analyses were conducted for boys and girls separately. The  
22 associations between normalized handgrip strength in kg/body mass in kg and four  
23 behaviours (smoking status [number of cigarettes smoked per day], body mass index, CRF

1 [shuttle-runs], and diet [score ranges from -4 to 12 points]) and three factors (total  
2 cholesterol, blood pressure and glucose) were tested by means of *Pearson* correlation  
3 coefficients. All analyses were performed using the Statistical Package for Social Sciences  
4 (v. 22.0 for WINDOWS, Chicago, USA), and the level of significance was set to 0.05.

5

## 6 **RESULTS**

7 The 1,199 scholars included 627 boys, and mean age was 13.1 (2.2) years. Boys had  
8 lower levels of total cholesterol than girls ( $p < 0.001$ ), and girls had lower cardiorespiratory  
9 fitness, handgrip strength and normalized handgrip strength ( $p < 0.001$ ) (Table I).

10 **\*\*\*Table I about here\*\*\***

11 Higher levels of handgrip and normalized handgrip strength were associated with a  
12 higher number of ideal CVH components in both boys ( $p$  for trend  $< 0.001$ ) and girls ( $p$  for  
13 trend  $< 0.001$ ) (Figure 1).

14 **\*\*\*Figure 1 about here\*\*\***

15 Higher levels of handgrip strength were associated with a higher number of ideal  
16 health behaviours ( $p$  for trend  $< 0.001$  in both boys and girls) (Figure 2A and Figure 2C),  
17 and with a higher number of ideal health factors in boys ( $p$  for trend  $< 0.001$ ) (Figure 2B  
18 and Figure 2D).

19 **\*\*\*Figure 2 about here\*\*\***

20 Levels of handgrip strength (both handgrip and normalized handgrip strength) were  
21 different between ideal versus non-ideal components except for glucose or total cholesterol  
22 groups in girls (Table II). Overall, similar results were observed when we included physical  
23 activity instead of CRF (data not shown). Finally, in both sexes, we found an inverse  
24 correlation between normalized grip strength and number cigarettes smoked per day ( $r = -$

1 0.356,  $P < 0.01$ ), BMI ( $r = -0.604$ ,  $P < 0.01$ ), CRF ( $r = -0.424$ ,  $P < 0.01$ ), diet score ( $r = -$   
2  $0.104$ ,  $P = 0.45$ ), total cholesterol ( $r = -0.238$ ,  $P < 0.01$ ), blood pressure ( $r = -0.220$ ,  $P < 0.05$ )  
3 and glucose ( $r = -0.016$ ,  $P < 0.01$ ).

4 **\*\*\*Table II about here\*\*\***

## 5 **DISCUSSION**

6 The findings of the present study indicate that handgrip strength is positively  
7 associated with the ideal CVH index among Colombian children and adolescents. The  
8 importance of muscular strength is recognized in most current recommendations for  
9 maintaining and improving health status, and preventing chronic diseases<sup>45</sup>. A recent meta-  
10 analysis highlights the importance of developing muscular strength in youth for a number  
11 of health-related benefits in young population<sup>46</sup>. Also, our study suggests a positive link  
12 between ideal CVH metrics and handgrip (for both absolute and relative values). In spite of  
13 the fact that there are very few studies on this topic, Ruiz et al.<sup>17</sup> showed that higher levels  
14 of CRF were associated with a higher number of ideal CVH components in both boys and  
15 girls. These findings together with our results confirm that physical fitness should be  
16 considered a hallmark factor for meeting ideal CVH components.

17 Several studies have showed the relationship between individual components  
18 included in the ideal CVH and handgrip strength in children and adolescents. As the  
19 American College of Sports Medicine recently recommended the incorporation of grip  
20 strength testing as a component of musculoskeletal fitness assessment in children<sup>47</sup>, it is  
21 important to not only understand the link between variability in this measured outcome and  
22 that of health risks, but also from the context of translating meaningful risk-stratification  
23 information to clinical and public health audiences. Moreover, a very recent systematic

1 review and meta-analysis revealed strong evidence for an inverse association between  
2 musculoskeletal fitness and cardiometabolic risk factors among adolescents<sup>32</sup>.

3       Regarding health behaviours, there is a well-established link between BMI, physical  
4 activity, cardiorespiratory fitness, and muscular strength<sup>48</sup>. However, there is less evidence  
5 pertaining to the relationship between handgrip strength and healthy dietary adherence in  
6 younger populations. Therefore, considering the limited number of studies that have  
7 previously examined these associations, it is difficult to compare our results. A cross-  
8 sectional study in Spanish adolescents showed that there was not relationship between  
9 handgrip strength and ideal diet; the authors hypothesized that results seemed to be more  
10 associated with physical activity levels and aerobic capacity at these ages<sup>49</sup>. Regarding  
11 smoking habits, the small number of available studies has shown inconsistent results. A  
12 previous study reported that smoking habits were not related with handgrip strength<sup>49</sup>;  
13 however, another Spanish study in children and adolescents suggests that those who had  
14 muscle weakness had a significantly higher odds ratio of reporting smoking tobacco  
15 sometimes<sup>50</sup>. In our study, we found an inverse correlation between normalized grip  
16 strength and number cigarettes smoked per day ( $r=-0.356$ ,  $P < 0.01$ ). Cross-sectional and  
17 longitudinal studies showed that smoking affects the body through, for example, increased  
18 oxidative stress, which negatively influences the muscles<sup>3,5,51</sup>. Circulating cigarette smoke  
19 constituents seem to play an important role in the underlying molecular mechanisms of  
20 muscle damage, such as reduced oxygen delivery and impair mitochondrial function<sup>52</sup>.

21       These findings are particularly important from a public health perspective, given the  
22 well-known negative consequences of smoking and the fact that this behavior starts already  
23 at young ages. Regarding health factors, the role of muscular strength in prevention of  
24 cardiovascular disease has become increasingly recognized<sup>39</sup>. Our results show higher

1 values of handgrip strength in children and adolescents who had ideal health factors as  
2 compared to peers that not meet ideal condition. The exception for this was for total  
3 cholesterol and glucose in girls, which could be explained by the small number of youth  
4 with non-ideal factors (2% and 6%, respectively).

5         The present findings lend strong support to the growing body of literature revealing  
6 a link between muscle weakness and increased cardiometabolic risk factors;<sup>7-10</sup> and yet, the  
7 mechanisms underlying this association are still to be determined. It has been hypothesized  
8 that one possible mechanism by which healthy muscular strength exerts favorable health  
9 effects may be its capacity to reduce chronic low-grade inflammation. In the Pan-American  
10 HELENA study, Artero et al.<sup>9</sup> found an inverse association between muscular strength with  
11 lower levels of markers of chronic inflammation such as C-reactive protein, complement  
12 factors C3 and C4, leptin and white blood cell counts in adolescents, even after adjusting  
13 for gender, age, cardiorespiratory fitness, maturation, and socioeconomic status. In  
14 addition, Steene-Johannessen et al.<sup>7</sup>, Cohen et al.<sup>8</sup> and Peterson et al.<sup>53</sup> reported strong  
15 evidence of the inverse associations between muscular strength and cardiometabolic risk  
16 factors such as HOMA index, triglycerides, and blood pressure, positive associations with  
17 markers of endothelial function, and lower arterial stiffness. Differences in body  
18 distribution of excess adiposity could be another explanation, as there are studies that  
19 suggest that muscular strength is a stronger influence on cardiometabolic  
20 abnormalities<sup>17,19,31,33</sup>. However, further research is needed to confirm these mechanisms,  
21 especially in the pediatric population.

22         There are some limitations to this study. The observations of our study are limited  
23 by the descriptive and cross-sectional design, and therefore direction of causality cannot be  
24 determined. Another limitation was that adherence to the Mediterranean diet was measured

1 by a self-administered questionnaire, so some of the questions may have been  
2 misinterpreted deliberately or unintentionally by some participants. Future research is  
3 needed to better describe the age- and sex-specific trajectories of strength as a predictor of  
4 comorbidities across the lifespan and, perhaps just as importantly, to apply robust analyses  
5 that can compartmentalize risk into hierarchical categories<sup>53</sup>. Finally, it should also be  
6 noted that the formation of the ideal CVH metrics relies on the use of binary variables and  
7 on the assumption that all health behaviours and factors contained in this index contribute  
8 the same to the final score<sup>17</sup>.

9         The findings of this study indicate that handgrip strength is positively associated  
10 with ideal CVH metrics in Colombian youths. These results provide an important public  
11 health message that children and adolescents do not necessarily have to reach all 7 metrics  
12 to gain CVH benefits. Moreover, the data suggest that preventive efforts should be focused  
13 on those with few ideal health behaviors or factors, and should target early development of  
14 handgrip strength to reduce the risk of premature health problems.

#### 15 **ABBREVIATIONS**

16 CVD: Cardiovascular disease

17 CVH: Cardiovascular health

18 CCI: Intraclass correlation coefficient

19 BMI: Body mass index

20 IOTF: International Obesity Task Force

21 CRF: Cardiorespiratory fitness

22



1    **ACKNOWLEDGEMENTS**

2            We thank the children and adolescents who participated in the study and their  
3    parents and teachers for their collaboration. We also acknowledge the members involved in  
4    fieldwork for their efforts, particularly physical activity and health masters students for  
5    their work in the Field-based fitness assessment.

6    **FIGURE LEGEND**

7    Figure 1. Association between handgrip strength and normalized grip strength [measured as  
8    grip strength in kg/body mass in kg] across ideal CVH metrics in schoolchildren. (A) Boys  
9    and (B) Girls.

10   Figure 2. Levels of handgrip strength and normalized grip strength [measured as grip  
11   strength in kg/body mass in kg] across ideal CVH behaviours (smoking, body mass index,  
12   cardiorespiratory fitness, and Mediterranean diet adherence) and ideal health factors (total  
13   cholesterol, blood pressure, and plasma glucose) in schoolchildren; (A-B) Boys and (C-D)  
14   Girls.

15   **REFERENCES**

- 16            1. Peterson MD, Krishnan C. Growth Charts for Muscular Strength Capacity With  
17            Quantile Regression. *Am J Prev Med* 2015; 49:935-8.
- 18            2. Ortega FB, Ruiz JR, Castillo MJ, Sjostrom M. Physical fitness in childhood and  
19            adolescence: a powerful marker of health. *Int J Obes (Lond)* 2008; 32:1-11.
- 20            3. Ortega FB, Silventoinen K, Tynelius P, Rasmussen F. Muscular strength in male  
21            adolescents and premature death: cohort study of one million participants. *BMJ*  
22            2012; 20;345:e7279.

- 1 4. Ruiz JR, Sui X, Lobelo F, Morrow JR Jr, Jackson AW, Sjöström M, Blair SN.  
2 Association between muscular strength and mortality in men; prospective cohort  
3 study. *BMJ* 2008; 337:a439.
- 4 5. Silventoinen K, Magnusson PK, Tynelius P, Batty GD, Rasmussen F. Association  
5 of body size and muscle strength with incidence of coronary heart disease and  
6 cerebrovascular diseases: A population-based cohort study of one million Swedish  
7 men. *Int J Epidemiol* 2009; 38:110-8.
- 8 6. Ruiz JR, Castro-Piñero J, España-Romero V, Artero EG, Ortega FB, Cuenca MM,  
9 et al. Field-based fitness assessment in young people: the ALPHA health-related  
10 fitness test battery for children and adolescents. *Br J Sports Med* 2011; 45:518-24.
- 11 7. Steene-Johannessen J, Anderssen SA, Kolle E, Andersen LB. Low muscle fitness is  
12 associated with metabolic risk in youth. *Med Sci Sports Exerc* 2009; 41:1361-7.
- 13 8. Cohen DD, Gómez-Arbeláez D, Camacho PA, Pinzon S, Hormiga C, Trejos-Suarez  
14 J, et al. Low Muscle Strength Is Associated with Metabolic Risk Factors in  
15 Colombian Children: The ACFIES Study. *PLoS ONE* 2014; 9:e93150.
- 16 9. Artero EG, Ruiz JR, Ortega FB, España-Romero V, Vicente-Rodríguez G, Molnar  
17 D, Gottrand F, et al. Muscular and cardiorespiratory fitness are independently  
18 associated with metabolic risk in adolescents: the HELENA study. *Pediatr Diabetes*  
19 2011; 12:704-12.
- 20 10. Magnussen CG, Schmidt MD, Dwyer T, Venn A. Muscular fitness and clustered  
21 cardiovascular disease risk in Australian youth. *Eur J Appl Physiol* 2012; 112:3167-  
22 71.
- 23 11. Berenson GS, Srinivasan SR, Bao W, Newman WP 3rd, Tracy RE, Wattigney WA.  
24 Association between multiple cardiovascular risk factors and atherosclerosis in

- 1 children and young adults. The Bogalusa Heart Study. *N Engl J Med* 1998;  
2 338:1650-6.
- 3 12. Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van HL, et al.  
4 Defining and setting national goals for cardiovascular health promotion and disease  
5 reduction: the American Heart Association's strategic impact goal through 2020 and  
6 beyond. *Circulation* 2010;121:586-613.
- 7 13. Shay CM, Ning H, Daniels SR, Rooks CR, Gidding SS, Lloyd-Jones DM. Status of  
8 cardiovascular health in US adolescents: prevalence estimates from the National  
9 Health and Nutrition Examination Surveys (NHANES) 2005-2010. *Circulation*  
10 2013;127:1369-76.
- 11 14. Shay CM, Ning H, Allen NB, Carnethon MR, Chiuve SE, Greenlund KJ, et al.  
12 Status of cardiovascular health in US adults: prevalence estimates from the National  
13 Health and Nutrition Examination Surveys (NHANES) 2003-2008. *Circulation*  
14 2012; 125:45-56.
- 15 15. Laitinen TT, Pahkala K, Magnussen CG, Viikari JS, Oikonen M, Taittonen L, et al.  
16 Ideal cardiovascular health in childhood and cardiometabolic outcomes in  
17 adulthood: the Cardiovascular Risk in Young Finns Study. *Circulation* 2012;  
18 125:1971-8.
- 19 16. Pahkala K, Hietalampi H, Laitinen TT, Viikari JS, Rönnekaa T, Niinikoski H, et al.  
20 Ideal cardiovascular health in adolescence: effect of lifestyle intervention and  
21 association with vascular intima-media thickness and elasticity (the Special Turku  
22 Coronary Risk Factor Intervention Project for Children [STRIP] study). *Circulation*  
23 2013; 127:2088-96.

- 1 17. Ruiz JR, Huybrechts I, Cuenca-García M, Artero EG, Labayen I, Meirhaeghe A, et  
2 al. Cardiorespiratory fitness and ideal cardiovascular health in European  
3 adolescents. *Heart* 2015; 101:766-73.
- 4 18. Younus A, Aneni EC, Spatz ES, Osondu CU, Roberson L, Ogunmoroti O, et al. A  
5 Systematic Review of the Prevalence and Outcomes of Ideal Cardiovascular Health  
6 in US and Non-US Populations. *Mayo Clin Proc* 2016; 91:649-70.
- 7 19. Isasi CR, Parrinello CM, Ayala GX, Delamater AM, Perreira KM, Daviglius ML, et  
8 al. Sex Differences in Cardiometabolic Risk Factors among Hispanic/Latino Youth.  
9 *J Pediatr* 2016 Jun 18. doi: 10.1016/j.jpeds.2016.05.037.
- 10 20. Garber MD, Sajuria M, Lobelo F. Geographical Variation in Health-Related  
11 Physical Fitness and Body Composition among Chilean 8th Graders: A Nationally  
12 Representative Cross-Sectional Study. *PLoS ONE* 2014; 9:e108053.
- 13 21. Rodríguez Valero FJ, Gualteros JA, Torres JA, Umbarila Espinosa LM, Ramírez-  
14 Velez R. Association between muscular fitness and physical health status among  
15 children and adolescents from Bogotá, Colombia. *Nutr Hosp* 2015; 32:1559-66.
- 16 22. González SA, Sarmiento OL, Cohen DD, Camargo DM, Correa JE, Páez DC, et al.  
17 Results from Colombia's 2014 Report Card on Physical Activity for Children and  
18 Youth. *J Phys Act Health* 2014; 11:S33-44.
- 19 23. Arsenault JE, Mora-Plazas M, Forero Y, Lopez-Arana S, Jáuregui G, Baylin A, et  
20 al. Micronutrient and anthropometric status indicators are associated with physical  
21 fitness in Colombian schoolchildren. *Br J Nutr* 2011 105:1832-42.
- 22 24. Gualteros JA, Torres JA, Umbarila-Espinosa LM, Rodríguez-Valero FJ, Ramírez-  
23 Vélez R. A lower cardiorespiratory fitness is associated to an unhealthy status

- 1 among children and adolescents from Bogotá, Colombia. *Endocrinol Nutr* 2015;  
2 62:437-46
- 3 25. Rodríguez-Bautista YP, Correa-Bautista JE, González-Jiménez E, Schmidt-  
4 RioValle J, Ramírez-Vélez R. Values of waist/hip ratio among children and  
5 adolescents from Bogotá, Colombia: The FUPRECOL Study. *Nutr Hosp* 2015;  
6 32:2054-61.
- 7 26. Ramírez-Vélez R, Morales O, Peña-Ibagon JC, Palacios-López A, Prieto-Benavides  
8 DH, Vivas A, et al. Normative Reference Values For Handgrip Strength In  
9 Colombian Schoolchildren: The Fuprecol Study. *J Strength Cond Res* 2016 Apr 21.  
10 doi: 10.1519/JSC.0000000000001459.
- 11 27. Ramírez-Vélez R, Rodrigues-Bezerra D, Correa-Bautista JE, Izquierdo M, Lobelo.  
12 Reliability of Health-Related Physical Fitness Tests among Colombian Children and  
13 Adolescents: The FUPRECOL Study. *PLoS ONE* 2015; 10:e0140875.
- 14 28. Committee on Fitness Measures and Health Outcomes in Youth, Food and Nutrition  
15 Board, Institute of Medicine; Pate R, Oria M, Pillsbury L, editors. Washington  
16 (DC). National Academies Press (US); 2012.
- 17 29. Ruiz JR, Castro-Pinero J, Espana-Romero V, Artero EG, Ortega FB, Cuenca MM,  
18 et al. Field-based fitness assessment in young people: the ALPHA health-related  
19 fitness test battery for children and adolescents. *Br J Sports Med* 2011; 45:518-24.
- 20 30. Ruiz JR, Castro-Pinero J, Artero EG, Ortega FB, Sjostrom M, Suni J, et al.  
21 Predictive validity of health-related fitness in youth: a systematic review. *Br J*  
22 *Sports Med* 2009; 43:909-23.
- 23 31. Artero E, Ruiz J, Ortega F, España-Romero V, Vicente-Rodríguez G, Molnar D, et  
24 al. Muscular and cardiorespiratory fitness are independently associated with

- 1 metabolic risk in adolescents: the HELENA study. *Pediatric Diabetes* 2011; 12:704-  
2 12.
- 3 32. Smith JJ, Eather N, Morgan PJ, Plotnikoff RC, Faigenbaum AD, Lubans DR. The  
4 health benefits of muscular fitness for children and adolescents: a systematic review  
5 and meta-analysis. *Sports Med* 2014; 44:1209-23.
- 6 33. Ortega FB, Ruiz JR, Castillo MJ, Moreno LA, González-Gross M, Wärnberg J, et  
7 al. Low level of physical fitness in Spanish adolescents. Relevance for future  
8 cardiovascular health (AVENA study). *Rev Esp Cardiol* 2005; 58:898-909.
- 9 34. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for  
10 child overweight and obesity worldwide: international survey. *BMJ* 2000;  
11 320:1240-3.
- 12 35. Zimmet P, Alberti KGMM, Kaufman F, Tajima N, Silink M, Arslanian S, et al. The  
13 metabolic syndrome in children and adolescents – an IDF consensus report. *Pediatr*  
14 *Diabetes* 2007; 8:299-306.
- 15 36. de Ferranti S, Gauvreau K, Ludwig DS, Neufeld EJ, Newburger JW, Rifai N.  
16 Prevalence of the metabolic syndrome in American adolescents: Findings from the  
17 Third National Health and Nutrition Examination Survey. *Circulation* 2004;  
18 110:2494-7.
- 19 37. Serra-Majem L, Ribas L, Ngo J, Ortega RM, García A, Pérez-Rodrigo C, et al.  
20 Food, youth and the Mediterranean diet in Spain. Development of KIDMED,  
21 Mediterranean Diet Quality Index in children and adolescents. *Public Health Nutr*  
22 2004; 7:931-5.
- 23 38. Ortega FB, Ruiz JR, Castillo MJ, Sjöström M. Physical fitness in childhood and  
24 adolescence: a powerful marker of health. *Int J Obes (Lond)*. 2008; 32:1-11.

- 1 39. Leger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run  
2 test for aerobic fitness. *J Sports Sci* 1988; 6:93-101.
- 3 40. Welk GJ, Laurson KR, Eisenmann JC, Cureton KJ. Development of Youth Aerobic-  
4 Capacity Standards Using Receiver Operating Characteristic Curves. *Am J Prev*  
5 *Med* 2011; 41:S111-6.
- 6 41. Lobelo F, Pate RR, Dowda M, Liese AD, Ruiz JR. Validity of cardiorespiratory  
7 fitness criterion-referenced standards for adolescents. *Med Sci Sports Exerc* 2009;  
8 41:1222-9.
- 9 42. Lobelo F, Pate RR, Dowda M, Liese AD, Daniels SR. Cardiorespiratory fitness and  
10 clustered cardiovascular disease risk in U.S. adolescents. *J Adolesc Health* 2010;  
11 47:352-9.
- 12 43. Tanner JM, Whitehouse RH. Clinical longitudinal standards for height, weight,  
13 height velocity, weight velocity, and stages of puberty. *Arch Dis Child* 1976;  
14 51:170-9.
- 15 44. Matsudo SMM, Matsudo VKR. Self-assessment and physician assessment of sexual  
16 maturation in Brazilian boys and girls – concordance and reproducibility. *Am J*  
17 *Hum Biol* 1994; 6:451-5.
- 18 45. Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin*  
19 *Nutr* 2006; 84:475-82.
- 20 46. Smith JJ, Eather N, Morgan PJ, Plotnikoff RC, Faigenbaum AD, Lubans DR. The  
21 health benefits of muscular fitness for children and adolescents: a systematic review  
22 and meta-analysis. *Sports Med* 2014; 44:1209-23.

- 1 47. Riebe D, Franklin BA, Thompson PD, Garber CE, Whitfield GP, Magal M, et al.  
2 Updating ACSM's Recommendations for Exercise Preparticipation Health  
3 Screening. *Med Sci Sports Exerc* 2015; 47:2473-9.
- 4 48. Galavíz KI, Tremblay MS, Colley R, Jáuregui E, López y Taylor J, Janssen I.  
5 Associations between physical activity, cardiorespiratory fitness, and obesity in  
6 Mexican children. *Salud Publica Mex* 2012; 54:463-9.
- 7 49. Grao-Cruces A, Fernández-Martínez A, Nuviala A. Association of fitness with life  
8 satisfaction, health risk behaviors, and adherence to the Mediterranean diet in  
9 Spanish adolescents. *J Strength Cond Res* 2014; 28:2164-72.
- 10 50. Padilla-Moledo C, Ruiz JR, Ortega FB, Mora J, Castro-Piñero J. Associations of  
11 muscular fitness with psychological positive health, health complaints, and health  
12 risk behaviors in Spanish children and adolescents. *J Strength Cond Res* 2012;  
13 26:167-73.
- 14 51. Kok MO, Hoekstra T, Twisk JW. The longitudinal relation between smoking and  
15 muscle strength in healthy adults. *Eur Addict Res* 2012;18:70-5.
- 16 52. Rom O, Kaisari S, Aizenbud D, Reznick AZ. Identification of possible cigarette  
17 smoke constituents responsible for muscle catabolism. *J Muscle Res Cell Motil*  
18 2012; 33:199-208.
- 19 53. Peterson MD, Zhang P, Saltarelli WA, Visich PS, Gordon PM. Low Muscle  
20 Strength Thresholds for the Detection of Cardiometabolic Risk in Adolescents. *Am*  
21 *J Prev Med* 2016; 50:593-9.



Table I. Characteristics of children and adolescents in Bogota, Colombia [mean (SD) or frequencies], by sex.

	Girls (n=572)	Boys (n=627)	All (n=1,199)
Age (years)	12.9 (2.2)	13.3 (2.2)	13.1 (2.2)
Body mass (kg)	42.1 (9.0)	45.8 (11.3)	44.0 (10.4)
Height (cm)	149.3 (10.2)	155.7 (13.7)*	152.6 (12.5)
Body mass index (kg/m <sup>2</sup> )	18.6 (2.3)	18.5 (2.0)	18.6 (2.2)
Tanner stage Prepuber/Puber/Postpuber, (%)	7/89/4	6/84/10	6/86/8
Resting blood pressure			
Systolic blood pressure (mm Hg)	106.4 (11.1)	109.6 (12.4)	108.0 (11.9)
Diastolic blood pressure (mm Hg)	66.3 (7.7)	66.0 (8.0)	66.1 (7.8)
Glucose (mg/dL)	78.3 (13.3)	79.7 (12.9)	79.1 (13.1)
Total cholesterol (mg/dL)	140.9 (21.9)	133.3 (24.6)*	136.9 (23.6)
Mediterranean diet adherence (-4 to 12 points)	7.1 (2.0)	7.0 (1.8)	7.0 (2.1)
Cardiorespiratory fitness (mL/kg/min)	41.1 (3.9)	45.6 (4.7)*	43.5 (4.9)
Muscular strength			
Handgrip strength (kg)	19.6 (5.4)	25.3 (9.2)*	22.6 (8.2)
Normalized grip strength <sup>†</sup>	0.47 (0.08)	0.54 (0.11)*	0.51 (0.10)

<sup>†</sup> Handgrip strength/body mass.

\* *t*-test was applied to compare unadjusted means by sex ( $p < 0.001$ ).

Table II

Table II. Handgrip (kg) and normalized grip strength (measured as handgrip strength/body mass) mean and SE estimates by ideal CVH metrics, by sex.

	Girls (n=572)							Boys (n=627)								
	Ideal	Mean (SE)		Non-ideal	Mean (SE)		<i>p</i> value*	Ideal	Mean (SE)		Non-ideal	Mean (SE)		<i>p</i> value*		
	N/%	HG	HG/BM	N/%	HG	HG/BM		N/%	HG	HG/BM	N/%	HG	HG/BM			
Health behaviours							HG	HG/BM						HG	HG/BM	
Smoking	217/38	19.6 (0.3)	0.461 (0.004)	355/62	18.7 (0.4)	0.472 (0.005)	0.001	0.001	307/49	25.8 (0.5)	0.534 (0.006)	320/51	24.7 (0.5)	0.554 (0.006)	0.001	0.001
Body mass index	538/94	19.6 (0.2)	0.469 (0.003)	34/6	19.7 (0.8)	0.397 (0.011)	0.123	0.022	609/97	25.3 (0.3)	0.546 (0.004)	18/3	22.9 (2.2)	0.449 (0.021)	0.001	0.001
Cardiorespiratory fitness	440/77	20.6 (0.4)	0.465 (0.006)	132/23	19.4 (0.3)	0.469 (0.003)	0.001	0.001	545/87	26.0 (0.9)	0.545 (0.011)	82/13	25.2 (0.4)	0.544 (0.004)	0.001	0.001
Mediterranean diet adherence	103/18	19.9 (0.2)	0.465 (0.004)	469/82	18.4 (0.5)	0.464 (0.008)	0.001	0.001	94/15	25.8 (0.4)	0.548 (0.005)	533/85	22.4 (0.9)	0.518 (0.011)	0.001	0.001
Health factors																
Blood pressure	532/93	19.7 (0.2)	0.465 (0.003)	40/7	18.0 (0.7)	0.460 (0.011)	0.001	0.001	577/92	25.2 (0.4)	0.545 (0.004)	50/8	23.9 (1.7)	0.513 (0.017)	0.001	0.001
Glucose	561/98	20.6 (1.8)	0.482 (0.027)	11/2	19.8 (0.2)	0.464 (0.003)	0.498	0.090	602/96	25.6 (0.4)	0.545 (0.004)	25/4	23.0 (1.9)	0.540 (0.019)	0.001	0.001
Total cholesterol	538/94	19.9 (0.2)	0.464 (0.003)	34/6	19.2 (0.9)	0.471 (0.013)	0.594	0.298	596/95	25.9 (0.4)	0.547 (0.004)	31/5	18.3 (1.1)	0.492 (0.016)	0.001	0.001

\*From analysis of covariance with age as covariate and Tanner stage as random factor



HG: Handgrip (kg); HG/BM: normalized grip strength (measured as handgrip strength/body mass), CVH: cardiovascular health

Table I. Definition of the Ideal Cardiovascular Health Metrics (&lt;20 Years of Age) as Defined by the American Heart Association and the Criteria Used in this Study.



	Ideal Metric, AHA Definition	Ideal Metric, Definition in this Study
<i>Health behaviors</i>		
Smoking	Never tried; never smoked whole cigarette	Never smoked a cigarette
Body mass index	<85th percentile	<85th percentile
Physical activity	≥60 min of moderate- or vigorous-intensity activity every day	Healthy CRF was defined by using either the cut-off by sex- and age (shuttle-runs or estimated VO <sub>2peak</sub> ) listed in the healthy fitness zone
Diet	4–5 components*:  Fruit and vegetables: ≥4.5 cups/d Fish: 2 or more 3.5-oz‡ servings/wk Fiber-rich whole grains: 3 or more 1-oz-equivalent servings/d Sodium: <1500 mg/d Sugar-sweetened beverages: ≤450 kcal (36 oz)/wk	Mediterranean diet quality  Participants who had at least ≥8 points were categorised as having an ideal healthy diet
<i>Health factors</i>		
Total cholesterol	<170 mg/dL (<4.40 mmol/L)	<170 mg/dL
Blood pressure	<90th percentile	<90th percentile
Plasma glucose	<100 mg/dL (<5.6 mmol/L)	<100 mg/dL

**Figure 1**  
[Click here to download Figure: Fig 1.eps](#)

**A**



**B**



**Figure 2**  
[Click here to download Figure: Fig 2.eps](#)

