

Addressing Disease-Related Malnutrition in Healthcare: A Latin American Perspective

Journal of Parenteral and Enteral
 Nutrition
 Volume 40 Number 3
 March 2016 319–325
 © 2015 The Author(s)
 DOI: 10.1177/0148607115581373
 jpen.sagepub.com
 hosted at
 online.sagepub.com



Maria Isabel Correia, MD, PhD¹; Refaat A. Hegazi, MD, PhD, MPH²;
 José Ignacio Diaz-Pizarro Graf, MD³; Gabriel Gomez-Morales, MD, MPH, MS⁴;
 Catalina Fuentes Gutiérrez, MD⁵; Maria Fernanda Goldin, MD⁶;
 Angela Navas, MD⁷; Olga Lucia Pinzón Espitia, PhD⁸;
 and Gilmária Millere Tavares, MD⁹

Abstract

Alarming high rates of disease-related malnutrition have persisted in hospitals of both emerging and industrialized nations over the past 2 decades, despite marked advances in medical care over this same interval. In Latin American hospitals, the numbers are particularly striking; disease-related malnutrition has been reported in nearly 50% of adult patients in Argentina, Brazil, Chile, Costa Rica, Cuba, Dominican Republic, Ecuador, Mexico, Panama, Paraguay, Peru, Puerto Rico, Venezuela, and Uruguay. The tolls of disease-related malnutrition are high in both human and financial terms—increased infectious complications, higher incidence of pressure ulcers, longer hospital stays, more frequent readmissions, greater costs of care, and increased risk of death. In an effort to draw attention to malnutrition in Latin American healthcare, a feedM.E. Latin American Study Group was formed to extend the reach and support the educational efforts of the feedM.E. Global Study Group. In this article, the feedM.E. Latin American Study Group shows that malnutrition incurs excessive costs to the healthcare systems, and the study group also presents evidence of how appropriate nutrition care can improve patients' clinical outcomes and lower healthcare costs. To achieve the benefits of nutrition for health throughout Latin America, the article presents feedM.E.'s simple and effective Nutrition Care Pathway in English and Spanish as a way to facilitate its use. (*JPEN J Parenter Enteral Nutr.* 2016;40:319-325)

Keywords

malnutrition; nutrition; hospital; community; screening; assessment; oral nutrition supplement

All too often, patients are malnourished when they are admitted to hospitals around the world^{1,2} and likewise in Latin America.^{3,4} Alarming high rates of disease-related malnutrition have persisted in hospitals of both emerging and industrialized nations over the past 2 decades, despite marked advances in medical care over this same interval.^{3,5–12} In Latin American hospitals, the malnutrition numbers are particularly striking; disease-related malnutrition has been reported in nearly 50% of adult patients in Argentina, Brazil, Chile, Costa Rica, Cuba, Dominican Republic, Ecuador, Mexico, Panama, Paraguay, Peru, Puerto Rico, Venezuela, and Uruguay.^{3,8,13–16} Risk of malnutrition is highest among older people—in Latin America and worldwide.^{17–20}

During hospitalization, patients often have experiences that further worsen their nutrition status (eg, traditional preparation for surgery, missed mealtimes for medical procedures, and nil per os [nothing by mouth] orders).²¹ As a result, weight loss and nutrient deficits can slow recovery and raise the risk for disability.^{1,21} Unfortunately, hospital malnutrition is often overlooked and undertreated.^{3,22,23} Clinicians do not consistently follow best nutrition practices because barriers—lack of awareness, time, money, and training—stand in the way.^{24–26} While some malnourished patients are undertreated because their condition is unrecognized by clinicians, other very ill hospital patients are difficult to treat. Patients may arrive at the

hospital with acute or chronic conditions complicated by a history of poor nutrition, which may require specialized

From the ¹Universidade Federal, de Minas Gerais, Brazil; ²Abbott Nutrition Research and Development, Columbus, Ohio; ³General Surgery and Clinical Nutrition Department, Hospital Ángeles Lomas & Anáhuac University, Mexico City, Mexico; ⁴Abbott Nutrition, Bogotá, Colombia; ⁵Hospital Clínico Fuerza Aérea de Chile, Gral. Dr Raúl Yazigi, Santiago, Chile; ⁶Hospital Italiano Buenos Aires, Buenos Aires, Argentina; ⁷Clinica Reina Sofia, Bogotá, Colombia; ⁸Universidad del Rosario-Méderi, Bogotá, Colombia; and ⁹Centro Integrado de Atenção à Saúde, Vitória, Espírito Santo, Brazil.

Financial disclosure: The feedM.E. Latin American Study Group received a grant from Abbott Nutrition to fund the Latin American manuscript project; this grant was used for operational activities and for funding the Buenos Aires meeting of the feedM.E. Latin American Study Group in October 2014. Members of the group did not receive payment for work on the document.

Supplemental material is available for this article.

Received for publication November 19, 2014; accepted for publication February 10, 2015.

This article originally appeared online on April 16, 2015.

Corresponding Author:

Maria Isabel Correia, MD, PhD, Av. Carandaí, Universidade Federal, de Minas Gerais, Brazil, 246 apt. 902, Belo Horizonte, MG, 30130-060, Brazil. Email: isabel_correia@uol.com.br

Table 1. Prevalence of Malnutrition in Latin America.

Study	Population	Prevalence of Malnutrition
Waltzberg et al (2001) ¹⁵	Cross-sectional, multicenter study of adult patients in Brazilian hospitals in the public healthcare system, n = 4000	48.1% by SGA
Correia and Campos (2003) ³	Cross-sectional, multicenter study of adult patients in hospitals of 13 Latin American countries, n = 9348	50.2% by SGA
Wyszynski et al (2003) ⁸	Study of adult patients in 38 hospitals of Argentina, n = 1000	47% by SGA
Baccaro and Sanchez (2009) ¹³	Study of adult male patients in the internal medicine service of an Argentina hospital, n = 152	48.7% by SGA
Lara-Pulido and Guevara-Cruz (2012) ¹⁷	Patients ≥65 years of age hospitalized in Mexico, n = 769	53.6% by MNA-SF
Veramendi-Espinoza et al (2013) ¹⁴	Cross-sectional study of adult patients in surgery and medicine wards of a Peruvian general hospital, n = 211	46.9%
Gallegos Espinosa et al (2014) ¹⁶	Cross-sectional, multicenter study of adult patients in hospitals of Ecuador, n = 5355	37.1% by SGA

MNA-SF, Mini Nutritional Assessment—short form, specifically used to detect nutrition status of people older than 65 years²⁹; SGA, Subjective Global Assessment for adult populations.²⁸

feeding ingredients, formulas, and techniques. In some cases, ideal therapies for each condition are not yet known.

feedM.E. is a medical education (M.E.) initiative developed by international nutrition experts. The feedM.E. Global Study Group sought to heighten awareness of malnutrition in hospitals and at other healthcare sites, to educate hospital staff and patients about nutrition's role in recovery from illness and injury, and to provide practical guidance for clinicians to take action and improve nutrition care.¹ A feedM.E. Latin American Study Group was recently formed to support and extend the international call to action for awareness and treatment of disease-related malnutrition. Study group participants were selected for their experience and expertise in clinical nutrition and for their representation of countries with hospital nutrition care programs. This Latin American feedM.E. initiative has been endorsed by the Federación Latinoamericana de Terapia Nutricional, Nutrición Clínica y Metabolismo (FELANPE), the principal organization for nutrition professionals in Latin America. The feedM.E. initiative follows the nutrition practices and principles put forth in the *Declaration of Cancun*, known as *Declaración Internacional Sobre el Derecho a la Nutrición en los Hospitales*, which was developed and published by FELANPE and partners in 2008.²⁷ The objectives of this declaration were to ensure that all patients admitted to public- and private-sector hospitals in Latin America receive complete, adequate, quality, and timely nutrition care; to ensure that this care is provided by health professionals who are trained to prevent the risks associated with hospital malnutrition; and to use nutrition to improve patients' quality of life and survival and also to reduce costs of hospitalization.

This article from the feedM.E. Latin American Study Group compiles evidence of the high prevalence of disease-related malnutrition in Latin American hospitals. It also shows that malnutrition incurs excessive costs to healthcare systems, and it shows that attention to nutrition can improve patients'

clinical outcomes and lower healthcare costs. As an online supplement to this article, we provide feedM.E.'s simple and effective Nutrition Care Pathway translated into Spanish to facilitate its use throughout Latin America. This Pathway can be followed as a guide to identify and treat patients at risk of malnutrition in the community, to diagnose and treat malnutrition in hospitalized patients, and to follow up with postdischarge nutrition care in the community.

Hospital Malnutrition in Latin America: Prevalent and Persistent

From prevalence studies conducted since 2000, results show that nearly 50% of all people admitted to hospitals in Latin America are malnourished (Table 1).^{3,8,13,15–17} Malnutrition is determined by use of validated and reliable assessment tools, such as the Subjective Global Assessment tool for all adult populations and the Mini Nutritional Assessment (MNA) for patients older than 65 years.^{28,29} While the reported prevalence of malnutrition may vary according to the criteria used (serum albumin or other blood tests, assessment tools, anthropometric measures) and the population studied (elderly, critically ill, country), the conclusions are disturbingly consistent. Disease-related malnutrition is highly prevalent throughout Latin America, and this high prevalence has changed very little over more than a decade.

While malnutrition prevalence is high, physician awareness is low; as a result, nutrition therapy is underprescribed. In a classic survey, one-third of hospitalized Brazilian patients were not fed for more than 3 days, only 19% of patients' medical records noted nutrition status, and only 10% of patients received oral nutrition or enteral nutrition (EN) by tube feeding.¹⁵ Similarly, a study of Argentinian hospital patients found that <40% of charts contained any nutrition information, and fewer than 15% of charts recorded usual and current

weight and height.⁸ In Ecuador, a large, recent study in public hospitals showed a 37% prevalence of malnutrition among admitted patients, yet Ecuador lacks policies, resources, and educational programs to identify and address disease-related malnutrition.¹⁶

Malnutrition Leads to Poor Clinical Outcomes and Higher Costs of Care

Disease-related malnutrition takes high tolls on patients and on healthcare systems, too; malnutrition is associated with increased risks for morbidity and mortality and higher costs of care.¹ Results of numerous clinical studies show that malnourished patients in the hospital are at increased risk for complications such as pressure ulcers, infections, falls, and death. In a Brazilian hospital survey, malnutrition was directly associated with greater frequency and severity of pressure ulcers; nearly 100% of patients with stage II or worse pressure ulcers were malnourished.³⁰ In another Brazilian study, malnourished patients had hospital stays 7 days longer and hospital episode costs up to 3-fold higher, and they were more than twice as likely to die compared with their adequately nourished peers.⁴ Malnourished Brazilian patients admitted to the intensive care unit (ICU) were twice as likely to be readmitted and 8 times more likely to die than were their nourished comparators.³¹ In Cuban hospitals, malnutrition was prevalent and associated with infections and longer length of hospital stay.³² Among Brazilian patients recovering from hip fracture, poor nutrition status predicted mortality.³³ Even in the community, undernutrition is a major risk factor for death in older Latin Americans.³⁴

Nutrition Care Is Associated With Improved Outcomes and Lower Costs

What is the evidence that attention to nutrition care has value? A major objection to implementing hospital nutrition programs is cost—the cost of training as well as the added costs of specialized nutrition such as oral nutrition supplements. To answer the important question—“Is the cost worth the expenditure?”—results of health economic studies are needed.

The global medical literature now reports abundant data on outcome benefits from nutrition interventions across many patient groups.¹ For example, nutrition intervention during hospitalization was correlated with significantly improved patients' strength, lower incidence of infectious complications and pressure ulcers, enhanced quality-of-life scores, and even lower risk of death.^{35–40} Global evidence likewise shows the positive association of nutrition interventions and health economic outcomes. Patients who received nutrition-focused care during hospitalization had shorter lengths of stay by 2–10 days^{36,41,42} and a significantly lower likelihood of readmission.^{36,39,43} A keystone study from the United States reported significantly lower costs by nearly \$5000 per hospital episode for patients who received oral nutrition supplements during




Figure 1. Create an institutional culture that values nutrition care.

their hospitalization compared with matched subjects with similar illness severity who did not receive oral nutrition supplements.⁴² A Brazilian study of digestive surgery patients compared those who received conventional in-hospital nutrition therapy (control group) with those who also received home-based nutrition therapy in the preoperative and posthospital discharge periods (study group). Patients in both groups achieved similar nutrition benefits, but those in the study group had significantly lower hospital episode costs, mostly due to needing fewer days of hospitalization.⁴⁴ Additional evidence associating nutrition intervention with improved health and financial outcomes has accumulated rapidly in the past year.^{42,45–48}

Need for Enhanced Nutrition Awareness and Training

The high prevalence of disease-related malnutrition in Latin America, paired with low awareness of what to do about disease-related malnutrition, represents an opportunity to improve nutrition care by increasing education and training. Healthcare administrators, clinical leaders and educators, and bedside clinicians all need to know and believe in the importance of nutrition in health care.^{1,26} In fact, 2 recent studies in Canadian hospitals found that nurses and doctors alike recognized benefits to nutrition care and wanted to increase their education and training on nutrition management.^{49,50}

To this end, healthcare system leaders need to create a culture that values nutrition by making good nutrition a part of their system's mission and goals (Figure 1).⁵¹ To build a culture of nutrition value, healthcare professionals must first understand current evidence-based nutrition guidelines, such as those to be newly

published by FELANPE in early 2015. Hospital leaders can develop their own policies and protocols to reflect nutrition practice guidelines, or they may prefer to adopt or adapt ready-made protocols and practice algorithms from published guidelines.

Next, staff training and education programs are essential to transfer policies and guidelines to everyday practice.²⁵ Many teaching-learning models are available to meet the unique needs and resources of each hospital. Hospital nutrition programs can be formal or informal, such as grand rounds presentations, in-service training classes, bedside instruction for small groups, one-on-one training sessions, computer-based learning, and visual reminders such as posters and checklists. The feedM.E. initiative includes resources such as an educational monograph on disease-related malnutrition, a clinical handbook, and teaching slide sets (www.nutritionmatters.com). The development of these materials was supported by Abbott Nutrition (Abbott Laboratories, Abbott Park, IL), as is the ongoing updating and maintenance of the website.

A New Nutrition Care Pathway: *Detectar, Nutrir y Vigilar*

Developed by the feedM.E. Global Study Group, the Nutrition Care Pathway is a simple and efficient strategy that can be tailored for use in various healthcare settings (Figure 2; also see online Supplemental Figure S1 in Spanish). With this pathway, feedM.E. guidance uses *screen*, *intervene*, and *supervene* as reminders to improve identification and treatment of malnutrition and to promote routine follow-up care.¹ In Spanish, such guidance can be stated as *detectar, nutrir y vigilar*. The Latin American feedM.E. Study Group recognizes the importance of engaging dietitians, nurses, and physicians to play complementary roles in achieving optimal nutrition care.^{52,53} We invite further studies to compare use of this Nutrition Care Pathway with other methods for screening, assessment, treatment, and monitoring of nutrition status as part of quality improvement initiatives across various hospital settings.

Detectar

The first step is to screen each patient's nutrition status on hospital admission or on initiation of care and use a validated tool to identify risk of malnutrition. The feedM.E. Nutrition Care Pathway begins with 2 questions from the Malnutrition Screening Tool (MST): Has your food intake decreased? and Have you lost weight recently?⁵⁴ These initial screening questions are simple enough to be used by health caregivers at all levels and even by patients themselves or their family members as a way to identify risk of malnutrition.

The appetite loss and weight loss questions are paired with a quick clinical judgment about whether the patient's illness or injury adds risk of malnutrition.⁵⁵⁻⁵⁷ A contemporary definition of malnutrition defines the condition as 3 clinical syndromes: (1) *starvation-related malnutrition*, undernutrition in the absence of




Figure 2. The Nutrition Care Pathway, reprinted with permission from Elsevier from Correia M, Hegazi R, Higashiguchi T, et al. Evidence-based recommendations for addressing malnutrition in healthcare: an updated strategy from the feedM.E. Global Study Group, *J Am Med Dir Assoc.* 2014;15: 544-550. †For individuals who can tolerate oral feeding.

inflammation; (2) *chronic disease-related malnutrition*, nutrition inadequacy associated with chronic conditions that cause mild to moderate inflammation; and (3) *acute disease- or injury-related malnutrition*, undernutrition associated with conditions that elicit marked inflammatory responses.⁵⁸ Chronic conditions, such as heart failure or kidney disease, have moderate inflammation as a disease component, which raises the risk of malnutrition. Severe acute health crises, such as surgery, burns, or sepsis, cause marked inflammation, which worsens nutrition status quickly.

If nutrition screening identifies risk for malnutrition, quick action is key to better health outcomes. The feedM.E. Study Group recommends that health professionals consider immediate intervention with nutrition advice for increased oral intake of regular or fortified food or early initiation of oral nutrition supplements. In addition, the study group recommends use of additional validated tools to make a malnutrition diagnosis and to assess the severity of nutrition shortfalls. There is no single measure or tool capable of adequately assessing the risk of poor outcomes in all hospital patients.⁵⁹ At present, many investigators are examining ways

to combine tools for more accurate identification of malnutrition in specific populations, such as using SGA, prealbumin, and anthropometrics for nondialyzed patients with kidney disease⁶⁰; including C-reactive protein (CRP), a measure of inflammatory status, for patients with cancer⁶¹; and adding physical function tests such as handgrip strength or 6-minute walking for older hospitalized patients.⁶² More studies are needed comparing different malnutrition assessment tools within single patient populations. In the interim, clinical judgment and the SGA tool are most frequently used to identify disease-related malnutrition,²⁸ and the MNA is an effective alternative for older people.^{20,63}

In Latin America, the SGA²⁸ and the MNA²⁰ tools are commonly used in practice. Such tools rely on clinical judgment, which includes awareness that malnutrition is possible even in people who are obese. In fact, obese individuals commonly experience sarcopenia (ie, loss of muscle mass, strength, and/or function), a condition that is managed with nutrition and exercise.⁶⁴ More studies are needed comparing different malnutrition assessment tools within single patient populations. In the interim, clinical judgment and the SGA tool are most frequently used to identify disease-related malnutrition,²⁸ and the MNA is an effective alternative for older people.^{20,63}

Nutrir. Based on the nutrition assessment, nutrition targets are set for energy, protein, and other specific nutrients. Other decisions involve what to feed and how much to feed: feed to target with regular food, fortified food (especially high in energy and protein, as needed), oral nutrition supplements (ONS), tube-fed EN, or parenteral nutrition, as needed. In some cases, more than one method can be used to reach nutrition targets (eg, food plus oral nutrition supplements). In all cases, prompt interventions are advised to preserve and to prevent worsening of nutrition status. Among patients who were malnourished and in the hospital, results of a 1-day food intake audit showed that >50% ate less than half of the food offered to them, thus emphasizing a need for intensified nutrition planning and care.²² It is also important to treat other underlying causes (eg, chewing or swallowing problems)⁶⁵⁻⁶⁷ and consequences of malnutrition (eg, inflammation, muscle and/or weight loss)⁶⁸⁻⁷¹ that are identified during screening or assessment. Treatments may include specific nutrition strategies (eg, increasing protein⁶⁸ or adding ingredients with anti-inflammatory or immune enhancing properties,⁶⁹⁻⁷¹ such as eicosapentaenoic acid [EPA] and glutamine) and nonnutritive procedures such as increased physical activity or mobility for surgical patients,⁷² those with cancer,^{73,74} or patients recovering from critical illness.⁶⁹⁻⁷¹

Vigilar. As the patient's disease status changes, continue to monitor nutrition status routinely with rescreening, reassessment, and feeding adjustments as needed. Routine rescreening and follow-up adjustments are necessary during hospitalization. Even in patients who were adequately

nourished at the time of hospital admission, periodic screening and further assessment, when indicated, are advised during hospitalization.

Furthermore, nutrition care does not end when a patient is released from the hospital. The final step of the Nutrition Care Pathway is follow-up with continuing attention to meeting nutrition needs. Without treatment, poor nutrition status on discharge predicts hospital readmission within 30 days.⁷⁵ New focus on postdischarge nutrition planning⁷⁶ can lower costly hospital readmissions,⁷⁷ improve quality of life for patients,^{38,39} and, in some cases, even reduce risk of death.³⁷ Effective nutrition care necessitates development of a post-discharge nutrition plan and use of strategies to ensure the plan is implemented. Results of a systematic review of 6 randomized controlled trials (surgical and medical patients of older age) showed that postdischarge nutrition care with use of ONS had a positive effect on nutrition intake (energy) and nutrition status (weight) in all trials.⁷⁸ The feedM.E. Latin American Study Group thus recommends continued efforts to prevent and treat malnutrition for patients who have been discharged from the hospital into long-term care centers or into the community.

Call to Action for Nutrition Care in Latin America

Throughout Latin America, nutrition therapy in hospitals and long-term care centers still has a long way to go from current practice to evidence-based best practice. We, the members of the feedM.E. Latin American Study Group, recognize that nutrition care improves patient outcomes and reduces health-care costs. We now call on healthcare professionals throughout Latin America to take action with “*detectar, nutrir y vigilar.*” The simple and efficient feedM.E. Nutrition Care Pathway recommends screening all patients on admission or at initiation of care, providing supportive nutrition when needed, and giving routine follow-up care with postdischarge nutrition planning, treatment, and monitoring.

To meet goals of nutrition adequacy, we also emphasize the importance of educating all healthcare professionals to provide timely and effective nutrition care—including training for physicians, nurses, food technicians, medical and nursing students, and medical residents. We note that optimal nutrition care engages an interdisciplinary team of health professionals who provide a continuum of patient care from the community to the hospital, then back to a rehabilitation center or to the community again.

Acknowledgments

The feedM.E. Latin American Study Group thanks Cecilia Hofmann, PhD, for her valued assistance with compilation of the medical literature and with editing this English-language review article; her work was funded by Abbott Nutrition.

Statement of Authorship

All authors contributed to the conception and design of the research as well as to the acquisition, analysis, and interpretation of the data; drafted and critically revised the manuscript; agree to be fully accountable for ensuring the integrity and accuracy of the work; and read and approved the final manuscript.

References

- Correia M, Hegazi R, Higashiguchi T, et al. Evidence-based recommendations for addressing malnutrition in healthcare: an updated strategy from the feed.M.E. Global Study Group. *J Am Med Dir Assoc*. 2014;15:544-550.
- Norman K, Pichard C, Lochs H, Pirlich M. Prognostic impact of disease-related malnutrition. *Clin Nutr*. 2008;27(1):5-15.
- Correia MI, Campos AC. Prevalence of hospital malnutrition in Latin America: the multicenter ELAN study. *Nutrition*. 2003;19(10):823-825.
- Correia MI, Waitzberg DL. The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through a multivariate model analysis. *Clin Nutr*. 2003;22(3):235-239.
- Edington J, Boorman J, Durrant ER, et al. Prevalence of malnutrition on admission to four hospitals in England. The Malnutrition Prevalence Group. *Clin Nutr*. 2000;19(3):191-195.
- Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. *Clin Nutr*. 2003;22(4):415-421.
- Pirlich M, Schutz T, Kempes M, et al. Prevalence of malnutrition in hospitalized medical patients: impact of underlying disease. *Dig Dis*. 2003;21(3):245-251.
- Wyszynski DF, Perman M, Crivelli A. Prevalence of hospital malnutrition in Argentina: preliminary results of a population-based study. *Nutrition*. 2003;19(2):115-119.
- Chakravarty C, Hazarika B, Goswami L, Ramasubban S. Prevalence of malnutrition in a tertiary care hospital in India. *Indian J Crit Care Med*. 2013;17(3):170-173.
- Corkins MR, Guenter P, DiMaria-Ghalili RA, et al. Malnutrition diagnoses in hospitalized patients: United States, 2010. *JPEN J Parenter Enteral Nutr*. 2014;38(2):186-195.
- Karmakar PS, Pal J, Maitra S, et al. Prevalence of malnutrition and its correlation with various diseases in elderly patients in a tertiary care centre in eastern India. *J Indian Med Assoc*. 2010;108(11):754-756.
- Kirkland LL, Kashiwagi DT, Brantley S, Scheurer D, Varkey P. Nutrition in the hospitalized patient. *J Hosp Med*. 2013;8(1):52-58.
- Baccaro F, Sanchez A. Determination of hospital malnutrition: a comparison between the subjective global assessment and body mass index [in Spanish]. *Rev Gastroenterol Mex*. 2009;74(2):105-109.
- Veramendi-Espinoza LE, Zafra-Tanaka JH, Salazar-Saavedra O, et al. Prevalence and associated factors of hospital malnutrition in a general hospital; Peru, 2012 [in Spanish]. *Nutr Hosp*. 2013;28(4):1236-1243.
- Waitzberg DL, Caiaffa WT, Correia MI. Hospital malnutrition: the Brazilian national survey (IBRANUTRI): a study of 4000 patients. *Nutrition*. 2001;17(7-8):573-580.
- Gallegos Espinosa S, Nicolalde Cifuentes M, Santana Porben S. State of malnutrition in hospitals of Ecuador. *Nutr Hosp*. 2014;30(2):425-435.
- Lara-Pulido A, Guevara-Cruz M. Malnutrition and associated factors in elderly hospitalized. *Nutr Hosp*. 2012;27(2):652-655.
- Perez Cruz E, Lizarraga Sanchez DC, Martinez Esteves Mdel R. Association between malnutrition and depression in elderly [in Spanish]. *Nutr Hosp*. 2014;29(4):901-906.
- Imoberdorf R, Meier R, Krebs P, et al. Prevalence of undernutrition on admission to Swiss hospitals. *Clin Nutr*. 2010;29(1):38-41.
- Kaiser MJ, Bauer JM, Ramsch C, et al. Frequency of malnutrition in older adults: a multinational perspective using the Mini Nutritional Assessment. *J Am Geriatr Soc*. 2010;58(9):1734-1738.
- Krumholz HM. Post-hospital syndrome—an acquired, transient condition of generalized risk. *N Engl J Med*. 2013;368(2):100-102.
- Agarwal E, Ferguson M, Banks M, Bauer J, Capra S, Isenring E. Nutritional status and dietary intake of acute care patients: results from the Nutrition Care Day Survey 2010. *Clin Nutr*. 2012;31(1):41-47.
- Schindler K, Pernicka E, Laviano A, et al. How nutritional risk is assessed and managed in European hospitals: a survey of 21,007 patients findings from the 2007-2008 cross-sectional nutritionDay survey. *Clin Nutr*. 2010;29(5):552-559.
- Cahill NE, Murch L, Cook D, Heyland DK. Barriers to feeding critically ill patients: a multicenter survey of critical care nurses. *J Crit Care*. 2012;27(6):727-734.
- Jones NE, Suurdt J, Ouellette-Kuntz H, Heyland DK. Implementation of the Canadian Clinical Practice Guidelines for Nutrition Support: a multiple case study of barriers and enablers. *Nutr Clin Pract*. 2007;22(4):449-457.
- Tappenden KA, Quatrara B, Parkhurst ML, Malone AM, Fanjiang G, Ziegler TR. Critical role of nutrition in improving quality of care: an interdisciplinary call to action to address adult hospital malnutrition. *JPEN J Parenter Enteral Nutr*. 2013;37(4):482-497.
- Comité Ejecutivo De La Declaracion. Declaración Internacional Sobre El Derecho A La Nutrición En Los Hospitales. http://www.felanpeweb.org/dec_cancun_2008.pdf. Accessed March 31, 2015.
- Detsky AS, McLaughlin JR, Baker JP, et al. What is subjective global assessment of nutritional status? *JPEN J Parenter Enteral Nutr*. 1987;11(1):8-13.
- Kaiser MJ, Bauer JM, Ramsch C, et al. Validation of the Mini Nutritional Assessment short-form (MNA-SF): a practical tool for identification of nutritional status. *J Nutr Health Aging*. 2009;13(9):782-788.
- Brito PA, de Vasconcelos Generoso S, Correia MI. Prevalence of pressure ulcers in hospitals in Brazil and association with nutritional status—a multicenter, cross-sectional study. *Nutrition*. 2013;29(4):646-649.
- Fontes D, Generoso Sde V, Toulson Davisson Correia MI. Subjective global assessment: a reliable nutritional assessment tool to predict outcomes in critically ill patients. *Clin Nutr*. 2014;33(2):291-295.
- Barreto Penie J; Cuban Group for the Study of Hospital M. State of malnutrition in Cuban hospitals. *Nutrition*. 2005;21(4):487-497.
- Gumieiro DN, Rafacho BP, Goncalves AF, et al. Mini Nutritional Assessment predicts gait status and mortality 6 months after hip fracture. *Br J Nutr*. 2013;109(9):1657-1661.
- Ferreira LS, do Amaral TF, Marucci Mde F, Nascimento LF, Lebrao ML, Duarte YA. Undernutrition as a major risk factor for death among older Brazilian adults in the community-dwelling setting: SABE survey. *Nutrition*. 2011;27(10):1017-1022.
- Barr J, Hecht M, Flavin KE, Khorana A, Gould MK. Outcomes in critically ill patients before and after the implementation of an evidence-based nutritional management protocol. *Chest*. 2004;125(4):1446-1457.
- Cawood AL, Elia M, Stratton RJ. Systematic review and meta-analysis of the effects of high protein oral nutritional supplements. *Ageing Res Rev*. 2012;11(2):278-296.
- Milne AC, Potter J, Vivanti A, Avenell A. Protein and energy supplementation in elderly people at risk from malnutrition. *Cochrane Database Syst Rev*. 2009;(2):CD003288.
- Norman K, Kirchner H, Freudenreich M, Ockenga J, Lochs H, Pirlich M. Three month intervention with protein and energy rich supplements improve muscle function and quality of life in malnourished patients with non-neoplastic gastrointestinal disease—a randomized controlled trial. *Clin Nutr*. 2008;27(1):48-56.
- Starke J, Schneider H, Altheld B, Stehle P, Meier R. Short-term individual nutritional care as part of routine clinical setting improves outcome and quality of life in malnourished medical patients. *Clin Nutr*. 2011;30(2):194-201.
- Stratton RJ, Ek AC, Engfer M, et al. Enteral nutritional support in prevention and treatment of pressure ulcers: a systematic review and meta-analysis. *Ageing Res Rev*. 2005;4(3):422-450.

41. Lee C, Rucinski J, Bernstein L. A systematized interdisciplinary nutritional care plan results in improved clinical outcomes. *Clin Biochem.* 2012;45:1145-1149.
42. Philipson TJ, Snider JT, Lakdawalla DN, Stryckman B, Goldman DP. Impact of oral nutritional supplementation on hospital outcomes. *Am J Manag Care.* 2013;19(2):121-128.
43. Stratton RJ, Hebuterne X, Elia M. A systematic review and meta-analysis of the impact of oral nutritional supplements on hospital readmissions. *Ageing Res Rev.* 24 2013;12:884-897.
44. Baxter YC, Dias MC, Maculevicius J, Ceconello I, Coteleng B, Waitzberg DL. Economic study in surgical patients of a new model of nutrition therapy integrating hospital and home vs the conventional hospital model. *JPEN J Parenter Enteral Nutr.* 2005;29(1)(suppl):S96-S105.
45. Hamdy O, Ernst FR, Baumer D, Mustad V, Partridge J, Hegazi R. Differences in resource utilization between patients with diabetes receiving glycemia-targeted specialized nutrition vs standard nutrition formulas in U.S. hospitals. *JPEN J Parenter Enteral Nutr.* 2014;38(2)(suppl):86S-91S.
46. Lakdawalla DN, Mascarenhas M, Jena AB, et al. Impact of oral nutrition supplements on hospital outcomes in pediatric patients. *JPEN J Parenter Enteral Nutr.* 2014;38(2)(suppl):42S-49S.
47. Linthicum MT, Thornton Snider J, Vaithianathan R, et al. Economic burden of disease-associated malnutrition in China [published online October 8, 2014]. *Asia Pac J Public Health.*
48. Thornton Snider J, Jena AB, Linthicum MT, et al. Effect of hospital use of oral nutritional supplementation on length of stay, hospital cost, and 30-day readmissions among Medicare patients with COPD [published online October 30, 2014]. *Chest.*
49. Duerksen DR, Keller HH, Vesnaver E, et al. Physicians' perceptions regarding the detection and management of malnutrition in Canadian hospitals: results of a Canadian Malnutrition Task Force Survey [published online June 2, 2014]. *JPEN J Parenter Enteral Nutr.*
50. Duerksen DR, Keller HH, Vesnaver E, et al. Nurses' perceptions regarding the prevalence, detection, and causes of malnutrition in Canadian hospitals: results of a Canadian Malnutrition Task Force Survey [published online September 4, 2014]. *JPEN J Parenter Enteral Nutr.*
51. Brantley SL. Implementation of the enteral nutrition practice recommendations. *Nutr Clin Pract.* 2009;24(3):335-343.
52. Jefferies D, Johnson M, Ravens J. Nurturing and nourishing: the nurses' role in nutritional care. *J Clin Nurs.* 2011;20(3-4):317-330.
53. Thoresen L, Rothenberg E, Beck AM, Irtun O. Doctors and nurses on wards with greater access to clinical dietitians have better focus on clinical nutrition. *J Hum Nutr Diet.* 2008;21(3):239-247.
54. Ferguson M, Capra S, Bauer J, Banks M. Development of a valid and reliable malnutrition screening tool for adult acute hospital patients. *Nutrition.* 1999;15(6):458-464.
55. Jensen GL, Hsiao PY, Wheeler D. Adult nutrition assessment tutorial. *JPEN J Parenter Enteral Nutr.* 2012;36(3):267-274.
56. Jensen GL, Mirtallo J, Compher C, et al. Adult starvation and disease-related malnutrition: a proposal for etiology-based diagnosis in the clinical practice setting from the International Consensus Guideline Committee. *JPEN J Parenter Enteral Nutr.* 2010;34(2):156-159.
57. White JV, Guenter P, Jensen G, Malone A, Schofield M. Consensus statement: Academy of Nutrition and Dietetics and American Society for Parenteral and Enteral Nutrition: characteristics recommended for the identification and documentation of adult malnutrition (undernutrition). *JPEN J Parenter Enteral Nutr.* 2012;36(3):275-283.
58. Jensen GL, Compher C, Sullivan DH, Mullin GE. Recognizing malnutrition in adults: definitions and characteristics, screening, assessment, and team approach. *JPEN J Parenter Enteral Nutr.* 2013;37:802-807.
59. van Bokhorst-de van der Schueren MA, Lonterman-Monach S, de Vries OJ, Danner SA, Kramer MH, Muller M. Prevalence and determinants for malnutrition in geriatric outpatients. *Clin Nutr.* 2013;32:1007-1011.
60. Ikizler TA, Cano NJ, Franch H, et al. Prevention and treatment of protein energy wasting in chronic kidney disease patients: a consensus statement by the International Society of Renal Nutrition and Metabolism. *Kidney Int.* 2013;84(6):1096-1107.
61. Tan CS, Read JA, Phan VH, Beale PJ, Peat JK, Clarke SJ. The relationship between nutritional status, inflammatory markers and survival in patients with advanced cancer: a prospective cohort study. *Support Care Cancer.* 2015;23(2):385-391.
62. Martín-Ponce E, Hernandez-Betancor I, Gonzalez-Reimers E, Hernandez-Luis R, Martinez-Riera A, Santolaria F. Prognostic value of physical function tests: hand grip strength and six-minute walking test in elderly hospitalized patients. *Sci Rep.* 2014;4:7530.
63. Guigoz Y, Lauque S, Vellas BJ. Identifying the elderly at risk for malnutrition. The Mini Nutritional Assessment. *Clin Geriatr Med.* 2002;18(4):737-757.
64. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing.* 2010;39(4):412-423.
65. Cabrera MA, Mesas AE, Garcia AR, de Andrade SM. Malnutrition and depression among community-dwelling elderly people. *J Am Med Dir Assoc.* 2007;8(9):582-584.
66. Loser C. Malnutrition in hospital: the clinical and economic implications. *Dtsch Arztebl Int.* 2010;107(51-52):911-917.
67. Morley JE. Weight loss in older persons: new therapeutic approaches. *Curr Pharm Des.* 2007;13(35):3637-3647.
68. Bauer J, Biolo G, Cederholm T, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. *J Am Med Dir Assoc.* 2013;14:542-559.
69. Braga M, Wischmeyer PE, Drover J, Heyland DK. Clinical evidence for pharmaconutrition in major elective surgery. *JPEN J Parenter Enteral Nutr.* 2013;37(5)(suppl):66S-72S.
70. Ellinger S. Micronutrients, arginine, and glutamine: does supplementation provide an efficient tool for prevention and treatment of different kinds of wounds? *Adv Wound Care (New Rochelle).* 2014;3(11):691-707.
71. Hegazi RA, Wischmeyer PE. Clinical review: optimizing enteral nutrition for critically ill patients—a simple data-driven formula. *Crit Care.* 2011;15(6):234.
72. Ljungqvist O. ERAS—enhanced recovery after surgery: moving evidence-based perioperative care to practice. *JPEN J Parenter Enteral Nutr.* 2014;38(5):559-566.
73. Fearon KC. Cancer cachexia: developing multimodal therapy for a multidimensional problem. *Eur J Cancer.* 2008;44(8):1124-1132.
74. Fearon K, Arends J, Baracos V. Understanding the mechanisms and treatment options in cancer cachexia. *Nat Rev Clin Oncol.* 2013;10(2):90-99.
75. Friedmann JM, Jensen GL, Smiciklas-Wright H, McCamish MA. Predicting early nonelective hospital readmission in nutritionally compromised older adults. *Am J Clin Nutr.* 1997;65(6):1714-1720.
76. White J, Stotts N, Jones S, Granieri E. Managing postacute malnutrition (undernutrition) risk. *JPEN J Parenter Enteral Nutr.* 2013;37:816-823.
77. Rosen B, Maddox P, Ray N. A position paper on how cost and quality reforms are changing healthcare in America: focus on nutrition. *JPEN J Parenter Enteral Nutr.* 2013;37:796-801.
78. Beck AM, Holst M, Rasmussen HH. Oral nutritional support of older (65 years+) medical and surgical patients after discharge from hospital: systematic review and meta-analysis of randomized controlled trials. *Clin Rehabil.* 2013;27(1):19-27.