



Metagenomic Analysis of Surface Waters and Wastewater in the Colombian Andean Highlands: Implications for Health and Disease

Vanessa Urrea¹ · Luisa Páez-Triana¹ · Natalia Velásquez-Ortiz¹ · Milena Camargo^{1,2} · Luz H. Patiño¹ · Laura Vega¹ · Nathalia Ballesteros¹ · Arsenio Hidalgo-Troya³ · Luis-Alejandro Galeano⁴ · Juan David Ramírez^{1,5} · Marina Muñoz^{1,6} 

Received: 16 June 2024 / Accepted: 2 December 2024 / Published online: 28 February 2025
© The Author(s) 2025

Abstract

Urban water bodies serve as critical reservoirs of microbial diversity, with major implications for public health and environmental quality. This study aimed to characterize the microbial diversity of surface waters and wastewater from the Pasto River in the Colombian Andean Highlands, offering insights that may support water quality monitoring efforts. Sampling was conducted at three river sites and one wastewater location. Standard physicochemical and microbiological analyses were performed, including real-time PCR to detect protozoan pathogens *Giardia* spp. and *Cryptosporidium* spp. Metagenomic sequencing provided an in-depth taxonomic and functional profile of microbial communities through two complementary approaches: (i) read-based analysis to identify abundant families and species, both pathogenic and beneficial, and (ii) detection of health-related molecular markers, including antimicrobial resistance markers and virulence factors. Physicochemical analyses showed distinct profiles between wastewater and surface water, with wastewater exhibiting elevated levels of suspended solids (113.6 mg/L), biochemical oxygen demand (BOD, 311.2 mg/L), and chemical oxygen demand (COD, 426.7 mg/L). Real-time PCR detected *Giardia* spp. DNA in 75% (76/102) of the samples and *Cryptosporidium* spp. DNA in 94% (96/102) of samples. The metagenomic read-based profiling identified *Aeromonas media* as a prevalent pathogen and *Polaromonas naphthalenivorans* as a potential biodegradative agent. The metagenomic assembly produced 270 high-quality genomes, revealing 16 bacterial species (e. g., *Acinetobacter johnsonii* and *Megamonas funiformis*) that provided insights into fecal contaminants and native aquatic microbes. Functional profiling further revealed a high prevalence of antimicrobial resistance markers, particularly for tetracyclines, aminoglycosides, and macrolides, with the highest abundance found in wastewater samples. Additionally, virulence factors were notably present in *Zoogloea ramigera*. The findings underscore the value of metagenomic profiling as a comprehensive tool for water quality monitoring, facilitating the detection of pathogens, beneficial species, and molecular markers indicative of potential health risks. This approach supports continuous monitoring efforts, offering actionable data for water management strategies to safeguard public health and maintain ecological integrity.

✉ Marina Muñoz
cmmunozd@unal.edu.co

¹ Centro de Investigaciones en Microbiología y Biotecnología -UR (CIMBIUR), Facultad de Ciencias Naturales, Universidad del Rosario, 110221 Bogotá, Colombia

² Centro de Tecnología en Salud (CETESA), Innovaseq SAS, 250027 Funza, Cundinamarca, Colombia

³ Grupo de Investigación Salud Pública, Departamento de Matemáticas y Estadística, Universidad de Nariño, 520002 Pasto, Colombia

⁴ Grupo de Investigación en Materiales Funcionales y Catálisis (GIMFC), Departamento de Química, Universidad de Nariño, 520002 Pasto, Colombia

⁵ Molecular Microbiology Laboratory, Department of Pathology, Molecular and Cell-Based Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

⁶ Instituto de Biotecnología-UN (IBUN), Universidad Nacional de Colombia, 111321 Bogotá, Colombia

Introduction

Water is a vital resource that covers approximately 71% of the Earth's surface. However, only 2.5% of this water is freshwater [1], essential for humans, animals, and ecosystems [2]. Surface waters refer to freshwater bodies on Earth's surface [3], including rivers, lakes, and reservoirs. They are critical for sustaining human development and providing numerous social and economic benefits to urban growth. The increased water demand, especially in urbanized and developing regions, has led to significant pressures on these water bodies [4]. Pollution from urbanization, agriculture, industry, and domestic activities, among other anthropogenic activities, has contributed to the degradation of water quality, particularly pronounced in developing countries [5] where resources for implementing water source conservation and treatment schemes are limited. This contamination has contributed to ecological harm and the spread of waterborne diseases that affect human and animal health, ultimately limiting the sustainable use of freshwater resources [6].

Growing concerns about water resources underscore the urgent need for applying appropriate tests to evaluate the state of freshwater and effective wastewater treatment strategies that address both traditional and emerging contaminants. Pollutants in water bodies often originate from diverse sources, including domestic, hospital, and wastewater discharge, frequently containing personal care products, pharmaceuticals, and other potentially harmful substances [7]. For that reason, monitoring the physicochemical parameters of surface waters provides valuable insights into the impact of human activity on water quality.

Water bodies host a diverse range of microorganisms, including protozoa, bacteria, and viruses, some of which are associated with fecal contamination [9] and antibiotic resistance [10], thereby posing additional risks. For example, protozoan pathogens such as *Giardia* spp. and *Cryptosporidium* spp. serve as indicators of fecal contamination and are commonly associated with waterborne intestinal diseases that can negatively impact human health [11], especially among vulnerable populations. However, beyond these pathogens, water bodies harbor complex microbial communities that require detailed characterization to better understand ecosystem processes [17], functionality, adaptability, and the services they provide [18]. Characterizing these communities helps assess potential risks to human health, animal welfare, and ecological integrity [8]. Integrating physicochemical and microbiological parameters is crucial for a comprehensive view, as aquatic environments can promote the persistence and spread of contaminants to multiple areas [9–16], highlighting the need for continuous monitoring and assessment.

The advent of next-generation sequencing (NGS) technologies has significantly advanced microbiome analyses of environmental samples. Particularly, metagenomics has emerged as a powerful tool for characterizing microbial communities, allowing for taxonomic profiling of diverse microorganisms, including those that are difficult or impossible to culture. For example, Chopyk et al. conducted a study in the United States that analyzed microbial composition in freshwater and saline surface waters over time. Beyond taxonomic classification, metagenomics enables the identification of antibiotic resistance genes, enhancing the understanding of the risks associated with using these water bodies in activities like agriculture [19]. Similarly, a study in Southern China profiled pathogenic microorganisms and antibiotic resistance genes in wastewater and surface waters, finding similar microbial compositions in both environments suggesting that pathogens in surface waters might be linked to the direct discharge of untreated wastewater [20]. In Korea, metagenomic-assembled genomes (MAGs) from surface water provided detailed information into microbial community composition, resistome, and species without known cultivated representatives [21]. A Brazilian study underscored the importance of monitoring surface waters for pathogens, including protozoa like *Cryptosporidium* spp. and *Giardia* spp. to strengthen surveillance efforts [22]. In Mexico, NGS enabled the taxonomic profiling of surface waters, facilitating the identification of pathogenic microorganisms and their associations with water quality [23]. Additionally, research in wastewater revealed molecular markers relevant to public health, such as antibiotic resistance genes, highlighting the role of sewage and its treatment plants in identifying these genetic markers [24].

In Colombia, drinking water quality risk is assessed using the IRCA (Water Quality Risk Index), which evaluates physicochemical and microbiological characteristics. Based on the results, Colombia states (departments) are classified as high, medium, low, or no risk regarding water quality. The department of Nariño, located in Southwest Colombia, is classified as having a medium level of water quality risk according to the IRCA, and reports indicate related health issues in the region, including waterborne diseases [25]. Pasto River is one of the primary rivers in Nariño's capital, supplies approximately 85% of the city's drinking water [26]. The river traverses three main basins with distinct uses: the upper basin supports agricultural activities, the middle basin flows through urban areas with significant domestic and industrial discharges, and the lower basin is predominantly used for irrigation [26–28]. Approximately, 90% of the city's domestic wastewater is discharged into the river without treatment, with the Juan XXIII collector contributing substantial pollution loads in the northern region [27]. Despite these potential risks, only traditional

microbiological methods have been used to assess the health hazards associated with the Pasto River.

The study aims to characterize the microbial communities in the surface and wastewater of the Pasto River, located in the Colombian Andean Highlands, to establish a baseline of microbial diversity and health-related molecular markers that can inform future monitoring and risk assessment strategies. Through traditional physicochemical analyses, real-time PCR detection of protozoan pathogens (*Giardia* spp. and *Cryptosporidium* spp.), and shotgun metagenomics, this research provides essential data for understanding the microbial composition and potential health risks associated with this water source. This study hypothesizes that surface waters and wastewater in the Pasto River contain diverse microbial communities with varying levels of pathogenicity and antimicrobial resistance, which could potentially impact public health, highlighting the need to employ these methods as a tool for monitoring water quality, supporting public health initiatives and environmental management strategies.

Materials and Methods

Sample Collection

This study was conducted on the Pasto River, the primary water resource for San Juan de Pasto, Colombia. Using EpiDat software (v.3.1) [29] for sample size calculation, we estimated the required samples based on an expected bacterial diversity of 79%, with a precision of 5% and a 95% confidence interval. This calculation was based on the research conducted by Köchling et al. [26], who collected samples from a river in north-eastern Brazil that was continuously exposed to industrial and domestic contaminants. These parameters required a minimum of five water samples for each sample type (surface water and wastewater). Over seven months from August 2022 to March 2023 (excluding January due to logistical issues), we collected 50 surface water samples from three locations—upper basin (S1), middle basin (S2), and lower basin (S3)—and 50 wastewater samples from the Juan XXIII wastewater collector (R4). This longitudinal sampling aimed to capture temporal shifts in microbiome composition across the four collection points.

Samples were collected using a plastic container attached to a 6-m rope, facilitating access to water bodies from bridges or the riverbank (Fig. S1-A). Before sampling, the container was rinsed with potable water and submerged multiple times in the surface water or wastewater to reduce contamination. For each point, two sampling methods were used: (i) single-sample collection, where 400 mL was collected at a specific time and location, and (ii) composite collection, which involved combining multiple sub-samples, obtaining 400 mL to represent basin-wide conditions. For

surface water samples, the collection was obtained every 15 min, while for wastewater, 12 sub-samples were collected hourly, to ensure the representation of the basin (Fig. S1-B).

Finally, 102 samples were gathered, comprising 16 from point S1, 20 from point S2, 19 from point S3, and 47 from point R4, totaling 47 wastewater samples and 55 surface water samples. To preserve sample integrity and reduce microbial changes before processing, the samples were stored in sterile glass containers and placed in a portable refrigerator, maintaining a temperature of approximately 4 °C.

Physicochemical Analysis of Surface and Wastewater Samples

Monthly aliquots of composite samples from S1 to R4 were analyzed for physicochemical parameters. *In-situ* measurements included pH, water and ambient temperatures, conductivity, and dissolved oxygen using a multiparameter instrument (HANNA instruments). Furthermore, turbidity, suspended solids, alkalinity, acidity, electrical conductivity, nitrites, chlorides, and phosphates were analyzed in a certified laboratory. Kruskal–Wallis test and Dunn's test with Benjamini–Hochberg adjustment were conducted to highlight the particular months where these significant differences occurred. For this, we include significant differences * (0.01–0.05) and very significant differences **, *** (0.001–0.01, <0.001).

Sample Preprocessing and DNA Extraction

In the laboratory, biomass from the samples was concentrated through serial filtration using 3.0, 0.45, and 0.22 µm pore-sized membranes (Millipore). Each 100 mL portion of the sample was sequentially filtered, and membranes were subsequently placed in sterile 50 mL Falcon tubes. The obtained biomass was collected by gently scraping the membrane with molecular-grade water in two rounds of 15 min each. After centrifugation, DNA was extracted from the resulting pellet using the Qiagen DNeasy Power Soil kit, following the manufacturer's guidelines, with the elution buffer prewarmed to 65 °C for optimal yield. DNA quality and quantity were assessed using the Nanodrop equipment (Thermo Fisher Scientific), and integrity was confirmed via agarose gel electrophoresis.

qPCR and Analysis for Protozoan Detection

Real-time PCR was employed to detect protozoan parasites *Giardia* spp. and *Cryptosporidium* spp. in water samples. The primer–probe sets used included GdF (5'-CATGCATGCCGCTCA-3'), GdR (5'-AGCGGTGTCCGGCTAGC-3'), and GdP (6FAM/AGGACAACGGTTGCAC/MGB) for

Giardia spp. with the following thermal profile: an initial 15-min phase at 50 °C, followed by 10 min at 95 °C. Subsequently, 40 cycles were carried out: 15 s at 95 °C, 1 min at 58 °C, and 1 min at 60 °C. CcF18S (5'-GTTTTTCATTAATCAAGAACGAAAGTTAGG-3'), CcR18S (5'-GAGTAA GGAACAACCTCCAATCTCTAG-3'), and CsP (6HEX/TCAGATACCGTCGTAGTCTTAACCATAAACTATGCC/TAMRA) primers and probes were employed for the detection of *Cryptosporidium* spp. [26], with a thermal profile of 95 °C for 10 min, followed by 45 cycles (modified to enhance detection sensitivity): 10 s at 95 °C and 30 s at 60 °C.

Metagenomic Sequencing and Data Analysis

Samples that met quality standards—average concentration > 40 ng/μL, 260/280 ratio between 1.8 and 2.0—were sent to Novogene for metagenomic sequencing. Targeting a minimum of 4 Gb raw reads per sample, sequencing was performed on an Illumina Novaseq platform using paired end reads of 150 bp.

After sequencing, reads were quality-checked using FastQC v.0.11.9 [30] and MultiQC v.1.6 [31]. Any adapter content, N content per base, or low-quality sequences were removed using Trimmomatic v.0.38 [32]. Taxonomic classification was performed using the Centrifuge software v.1.0.3-beta [33], with data visualization in Pavian [34] and RStudio. Additionally, analyses were conducted using eggNOG mapper v.2 [46], and metabolic pathways were elucidated using KofamKOALA [47] to explore functional aspects.

Assembly of Genomes from Metagenomes

High-quality reads were assembled into Metagenomic-Assemble Genomes (MAGs) using Metaspades v3.15.3 [37]. Binning was then performed with MetaBAT, Maxbin, and Concoct software [38–40], followed by refinement using DASTool [41]. The quality of the MAGs was assessed with CheckM v1.1.3 [42], ensuring completeness > 90% and contamination < 5%. Taxonomic assignment for the MAGs was conducted using GTDB-Tk v.1.7.0 [43], while genome annotation was performed with Prokka [44] and visualized with Proksee (<https://proksee.ca/>).

A pangenome analysis was conducted using the Roary [45], generating a core genome alignment that served as the basis for a phylogenetic reconstruction that was visualized in the interactive Tree of Life—iTOL (<https://itol.embl.de/>), to evaluate relationships among prominent microbiome members by comparison with publicly available reference genomes from BV-BRC, ensuring high completeness and quality in the comparisons.

Identification of Health-Related Molecular Markers

Two analytical approaches were applied to detect molecular markers associated with antibiotic resistance (AMR-MM) and virulence factors (VF). The first approach used high-quality reads analyzed with the Resistance Gene Identifier (RGI) software, while the second approach was based on MAGs, using ABRICATE software. In both cases, the annotation of these markers was performed by comparison with the CARD v3.1.3 [35] and Virulence Factor Database (VFDB) [36] databases, respectively.

Results

Monitoring Physicochemical Parameters in Surface and Wastewater Samples

The traditional physicochemical parameters were subjected to a Shapiro–Wilks, revealing that only conductivity, dissolved oxygen percentage, pH, and water temperature exhibited a normal distribution. Other factors, such as fluorides, sulfates, and suspended solids, did not conform to this regular distribution pattern. We used the Kruskal–Wallis test for non-parametric data, which showed no significant differences across months. An ANOVA test indicated that significant differences were present only for the pH variable for parametric data. A subsequent Dunn's test with Benjamini–Hochberg adjustment revealed notable variations in pH levels, specifically between March and November.

Despite the absence of statistically significant differences, a descriptive comparison was made between wastewater and surface water. Wastewater shows higher average water and ambient temperatures, with a difference of approximately 3.5 °C compared to surface water. Concerning phosphates, both wastewater (16.4 mg/L) and surface water (5.8 mg/L) samples exhibit elevated concentrations. In contrast, nitrites have a higher average value in surface water (0.2 mg/L) than in wastewater (0.1 mg/L). Suspended solids in wastewater (113.6 mg/L) exceed those in surface water (72 mg/L) by roughly 41.6 mg/L. Lastly, both BOD—Biochemical oxygen demand and COD—Chemical Oxygen Demand values are higher in wastewater (311.2 mg/L; 426.7 mg/L) in comparison to surface water (163.9; 269.4 mg/L) (Table S1).

Detection of *Giardia* spp. and *Cryptosporidium* spp. in Surface and Wastewater Samples

Regarding the microbiological component, the genetic material of *Giardia* spp. and *Cryptosporidium* spp. was identified using real-time PCR. These protozoan parasites have been responsible for widespread gastrointestinal diseases, sometimes leading to fatalities, especially among

children and individuals with weakened immune systems [48]. The DNA of *Giardia* spp. was detected in 76 out of 102 samples (75%), with 34 positive samples from wastewater and 42 from surface water. Samples without detectable *Giardia* spp. DNA was concentrated in three specific months. In November, six negative samples were obtained, two from the S3 surface water sampling point and four from the R4 wastewater point.

Another instance where *Giardia* spp. was not detected in specific samples collected in February. This month, 12 negative samples were distributed across all sampling points, with two at point S1, one at point S2, three at S3, and six at point R4. Finally, seven negative samples were identified in March, with two at S1, two at S3, and three at R4. Regarding detecting *Cryptosporidium* spp., it was found in 96 out of 102 samples (94%). Notably, the detection of *Cryptosporidium* spp. was higher throughout the study months, except for October (where one sample from the R4 wastewater point tested negative), November (with four negative samples at the R4 point), and March (with no detection at the S1 surface water sampling point).

Description of the Aquatic Microbiome: A Taxonomic Profiling from Reads

Metagenomic analysis was conducted over 74/102 (72.5%) samples that successfully passed the sequencing quality control checks. Sequences were not trimmed, as they exhibited high-quality with scores exceeding 34 in phred score result, with an average of 33.4 million reads per sample. The analysis employed the entire dataset (Tables S2, S3, S4). A total of 52.8% of the reads were taxonomically classified using Centrifuge software. The sample composition was predominantly microbial, with bacteria accounting for over 98%. Most of the reads that did not classify as Bacteria corresponded to *Homo sapiens*, with a higher relative abundance in wastewater (0.60) compared to surface water (0.39). The most abundant archaea species in surface water samples were *Methanobrevibacter smithii* and *Methanococcus maripaludis*. *Methanosarcina barkeri* was the dominant species in wastewater. Lastly, the most abundant assigned viruses globally in the samples were *Aeromonas virus phiO18P* and *Acanthamoeba polyphaga mimivirus*. Notably, no reads were associated with protozoa or fungi from the reads. In pursuit of these, additional analyses were conducted, entailing the extraction of the 18S-rRNA region and subsequent comparison with the Kraken2 database [49]. However, these accounted for less than 0.08% of the total, prompting the decision not to pursue these analyses due to potential inaccuracies in taxonomic assignment.

Temporal Analysis of Bacterial Family Dynamics in Wastewater and Surface Waters

We examined the prevalence of predominant bacterial families at each sample point along the San Juan de Pasto map. Figure 1 shows the locations: Point S1 is in the southern part of the city, Point S2 is in the central district, and Point S3 is in the northern region. The wastewater sampling point, R4, is also located in the north part of the city, upstream from Point S3, suggesting potential contamination from R4 point to S3 (Fig. 1). Comparisons were not performed for Site S1 due to the limited available samples ($N=1$). To investigate potential variations among microbial families across different sampling months, we conducted a Kruskal–Wallis analysis.

No significant differences (P -value > 0.05) were observed for Site S2. However, differences did emerge between Sites S3 and R4. A Dunnett test with Benjamini–Hochberg adjustment was conducted to highlight the specific months with significant differences. Site S3 showed significant differences for all families, as illustrated in Fig. S2. Variations within the Bacteroidaceae family were evident between February and September. Likewise, notable differences were identified in the Rhodocyclaceae and Pseudomonadaceae families between February and November (Fig. S2).

Moreover, variations in family composition were observed at wastewater point R4 across the months (Fig. S3). It is worth highlighting the significance of substantial differences, indicated by three asterisks (***) and characterized by P -value values below 0.001, as illustrated in Fig. S3. The Bacteroidaceae family exhibited significant differences, with a P -value of 0.002 when comparing September to February and 0.006 when comparing September to March. Additionally, variations were evident between October and February (P -value 0.009). The Burkholderiaceae family displayed variations in October compared to February (P -value 0.0006) and November (P -value 0.002). In contrast, the Campylobacteraceae family presented significant differences in February compared to August (P -value 0.00002) and December (P -value 0.00009). Similarly, the Comamonadaceae family showed variations between February and October (P -value 0.0002) and during August and February (P -value 0.001). The Enterobacteriaceae family revealed significant differences in March compared to October (P -value 0.0007) and between March and September (P -value 0.002). Concerning the Moraxellaceae family, changes in abundance were evident in October compared to December (P -value 0.008) and March (P -value 0.008). When examining changes in the Pseudomonadaceae family, significant differences were identified in November compared to February (P -value 0.0003) and October (P -value 0.00004). Lastly, the Shewanellaceae family, in addition to its distinctive profile relative to surface water points, exhibited variations in February and December




Fig. 1 The top 10 most abundant bacterial families at the sampling points in the Pasto River. *S1* wastewater sampling point 1, *S2* surface water sampling point 2, *S3* surface water sampling point 3, *R4* wastewater collector point

(P -value 0.00008) and between February and November (P -value 0.00001).

Abundant Bacterial Species with Pathogenic and Beneficial Traits

The top 20 most abundant bacterial species were identified across sites, resulting in 44 highly abundant species shared among them. Figure 2A presents all the bacteria for which there are reports of being potentially pathogenic. Particularly abundant species include *Bacteroides cellulosilyticus*, *Raoultella ornithinolytica*, and *Pseudomonas aeruginosa*. *Aeromonas media* was the most abundant species in this classification. On the other hand, although the *Burkholderia cepacia* complex is not the most abundant, its presence is noteworthy among the top 44. Otherwise, beneficial bacteria included *Thauera* sp. MZ1T and *Alicyciphilus denitrificans* emerge as notable species. Lastly, *Polaromonas*

naphthalenivorans was the most abundant bacterium in this classification (Fig. 2B).

Assembly and Taxonomic Assignment from Metagenomes

Clean reads yielded 270 high-quality Metagenome-Assembled Genomes from (MAGs), meeting the Bowers et al. criteria of 90% completeness and <5% contamination. Taxonomic assignment performed using GTDB-Tk allowed the acquisition of 175 MAGs classified at the species level, which were then compared to cross-references from PubMLST [50] to identify potential matches with known microorganism species. Reference genomes corresponding to each assigned species were visualized by comparison with publicly available genomes identified in BV-BRC: Bacterial and Viral Bioinformatics Resource Center [51] as a step to those genomes. Our selection criteria prioritized public genomes that were complete and high-quality. Species that




Fig. 2 Global overview of the most abundant bacteria at all sampling points. **A** Bacteria associated with reported infections, **B** Bacteria with beneficial potential

lacked available reference genomes were excluded from further analyses.

Among assigned genomes, 50 originated from wastewater and 19 from surface water, representing 16 bacterial species. Prominent species included *Acinetobacter johnsonii*, *Megaomonas funiformis* and *Phocaeicola vulgatus*.

COG Functional Analysis and Pangenome Insights

Clusters of Orthologous Groups (COGs) analysis provided insights into the metabolic and functional diversity within predominant MAGs (Fig. 3). This overview highlights the notably high frequency within the "Function unknown" category. Moreover, across all samples, there was a more extensive range of subcategories within the overarching

"Metabolism" category. For *A. johnsonii* displayed in the first panel of the figure, the COGs L and J from the information storage and processing category were the most prevalent. *P. vulgatus* also consistently presented COG M from the cellular process and signaling category. Moreover, *M. funiformis* shown in the third panel of the figure showed that within the COGs, the category "Metabolism" predominated over the others.

The genomes of the predominant MAGs species were analyzed alongside previously described reference genomes. For this analysis, we included MAGs with more than one bin and more than one assigned reference genome. The comparisons were visually represented using Proseek (Fig. 4A), and then midpoint-rooted trees for each species from core genome of each species were generated (Fig. 4B). Reference




Fig. 3 Clusters of Orthologous Groups (COGs) for the predominant species. **A** *Acinetobacter johnsonii*, **B** *Phocaecicola vulgatus*, and **C** *Megamonas funiformis*




Fig. 4 Phylogenetic reconstruction based on core genome alignment of the predominant Metagenome-assembled genomes. **A** *Acinetobacter johnsonii*, **B** *Phocaicola vulgatus*, and **C** *Megamonas funiformis*

genomes are marked in purple (to access these, please refer to the following link in Table S5). In contrast, the studied genomes are marked in blue. Regarding *Acinetobacter johnsonii*, all study genomes clustered within the same clade. Interestingly, the reference genome did not cluster with the other reference genomes or ours. For *Megamonas funiformis*, most study genomes formed a distinct cluster. Lastly, in the case of *Phocaeicola vulgatus*, two of our genomes were closely grouped. In contrast, the remaining genome (Sample DP46) was the most distant and distinct compared to the other genomes, including the reference genomes.

Relative Abundance of Molecular Markers of Health Interest from Reads

Molecular markers associated with antibiotic resistance were screened by comparison with the CARD database. Our research revealed the ten most frequently detected molecular markers related to drugs in wastewater and surface water samples. In the wastewater samples, indicated by dark green in the outer ring (R4), we found that the most common antibiotic resistance molecular markers, shown by their wide bands in the circus (Fig. 4A), were associated with tetracyclines, aminoglycosides, and macrolides. Conversely, in the surface water samples (S1–S3), marked in light blue on the outer ring, aminoglycosides and tetracycline consistently dominated. Additionally, sulfonamides were frequently observed at points S1 and S3. Reflecting the

trends in wastewater, macrolides were most prominent at point R4, following the other two drugs (Fig. 5A).

Moreover, we selected the top 15 molecular markers frequently detected in both wastewater and surface water samples. In the upper panel marked in pink in Fig. 5B, the markers associated with antibiotic resistance can be observed. Each marker's frequency is indicated, with a maximum frequency of 29 for surface waters and 42 for wastewater. This emphasizes the prevalence of markers *ermK* and *ermY* in surface waters compared to sewage. In contrast, *sul1* and *sul2* were more common in wastewater than in surface waters. Regarding the molecular markers associated with virulence factors, located in the lower panel marked in green, we found that markers *hlyA*, *mshE*, and *fliC* were more abundant in wastewater than in surface waters. Conversely, *gacS* and *sefC* were predominant in surface waters but absent in sewage.

While moving from the outermost to the innermost part of MAGs, the antibiotic resistance markers in red that align with CARD can be noticed. The reference genomes for each species can be found in purple, and in gray, the assembled genome from the study, along with its respective contigs, can be seen. Our study successfully identified various microbial species. For instance, we detected *Acinetobacter johnsonii* in six different genome bins. When we examined its genome, we found 939 contigs and three molecular markers in CARD. Similarly, we identified *Megamonas funiformis* in two genome bins, revealing a genome consisting of 505 contigs and one molecular marker in CARD.




Fig. 5 Molecular markers detected from Metagenome-assembled genomes. **A** Relationship between the frequency of genes coding for the most common drug-associated molecular markers at each sam-

pling point (S1–S3 surface water samples, R4 = wastewater samples). **B** Molecular markers associated with resistance (upper panel); molecular markers associated with virulence factors (lower panel)

Additionally, the genome of *Phocaeicola vulgatus*, identified in three bins, comprised 689 contigs and four molecular markers in CARD. Our study also revealed the presence of other species, including *Aliarcobacter cryaerophilus_A*, *Sphaerotilus montanus*, *Prevotella copri*, *Aeromonas media*, *Aeromonas rivipollensis*, *Bacteroides uniformis*, and *Phascolarctobacterium_A_succinatutens*. Furthermore, through analyses using high-quality WGS data, we assigned the following species: *Kaistella chaponensis*, *Trichococcus flocculiformis*, *Zoogloea ramigera*, *Alistipes putredinis*, *Lactococcus A raffinolactis*, and *Faecalibacterium prausnitzii*.

Also, we identified antimicrobial resistance markers. *Acinetobacter johnsonii*, for instance, exhibited 11 molecular markers associated with the beta-lactam resistance pathway, six genes linked to the Vancomycin resistance pathway, and five genes involved in the Cationic antimicrobial peptide (CAMP) pathway. Similarly, in *Megamonas funiformis*, we identified five genes within the beta-Lactam pathway, five in the Vancomycin pathway, and 5 in the CAMP pathway.

Molecular Markers of Interest in Health Associated with MAGs

Furthermore, using the previously obtained assemblages, we identified molecular markers related to health, specifically those associated with antibiotic resistance and virulence factors in the MAG species. We categorized resistance markers by species and their respective sampling points. This approach helped us identify the molecular markers most abundant within each species. In Fig. 6, genome assemblies for each species can be observed, depicted as circles. The color reflects the source of each assembly, indicating whether the sample originated from wastewater or surface

water and, in the latter scenario, the specific sampling point. The frequently detected antimicrobial resistance markers in wastewater (R4) included aminoglycoside, tetracycline, cephamycin, and cephalosporin, each found in three species.

In contrast, in surface waters (S3, S2), resistance markers for cephalosporins were identified in five species, while that for macrolide appeared in two. Furthermore, we identified virulence factors exclusively found in the species *Zoogloea ramigera*. In this case, we detected the genes *hsiC1/vipB*, *hsiC1/vipA*, and *pilT*.

Discussion

Studying the microbiome in aquatic environments is essential to understanding the microorganisms and their interactions, especially those that could impact public health [50]. From the findings of the physicochemical analyses, we can draw several conclusions. First, the elevated nitrite values in surface waters indicate contamination related to agricultural activities [51]. In contrast, the high values of temperature, suspended solids, and BOD in wastewater imply greater contamination, primarily due to domestic, hospital, and stormwater contaminants [52–54]. The identification of parasites like *Giardia* spp. and *Cryptosporidium* spp. aligns with findings from other studies. These parasites have been detected in various regions, including Spain in wastewater [55], Poland in surface waters [56], and within the Nariño (Colombia), as reported by Sánchez [61]. This underscores the concern associated with reusing surface water, as water serves to transmit these parasites [57]. These findings emphasize the potential public health risks and draw attention to the possible impact of the Juan XXIII wastewater




Fig. 6 Molecular health-related markers associated with the MAGs for each species. Each point represents an assembled genome of a specific species

collector, which discharges untreated water into surface waters.

Our study aimed to identify the most abundant families at each collection point to understand how they change over time. When examining the families present at the S3 surface water point (Figs. 1, S2), we noticed the widespread distribution of the Bacteroidaceae family in the environment. This is attributed to their large genomes, which enable them to adapt to diverse conditions [58]. Nevertheless, it is worth noting that some members of this family can act as opportunistic pathogens, potentially causing human infections. This suggests that there was likely a higher load of contaminants linked to mammalian excretions in September, as some of the bacterial species identified from these families thrive in the gastrointestinal system of particular mammal [59].

Similarly, we observed significant changes in the Rhodocyclaceae family. Members of this family have been identified in aquatic environments, as demonstrated by Chernitsyna et al. They also found that these family members play a role in bioremediation, particularly in denitrification processes. This aligns with our research, as we detected elevated levels of nitrites in surface waters, suggesting the possible involvement of these species in denitrification processes [60]. We also observed a similar pattern in the Pseudomonadaceae family. While some species are commonly found in the environment, it is essential to highlight the presence of pathogenic species that could pose public health concerns. The findings in surface waters provide valuable insights into identifying bacteria present in the environment, particularly emphasizing bacterial families that include members with pathogenic potential. This underscores the importance of monitoring this environment to comprehensively understand the associated risks when in contact with this valuable resource.

Regarding the families identified at wastewater sampling point R4 (Figs. 1, S3), we observed significant variations in the abundance of all families across different months. The Burkholderiaceae family exhibited higher abundance in September, suggesting increased contamination from hospital, domestic, and industrial wastewater sources [61]. We also noticed fluctuations in the Rhodocyclaceae family, which has been previously reported in wastewater treatment systems by Wang and colleagues. They identified some members of this family as participating in denitrification processes [62]. Another noteworthy finding was the higher abundance of the Campylobacteraceae family in February. This finding is significant as certain family members are associated with diarrheal diseases, indicating the presence of relevant pathogens [63].

Furthermore, identifying families such as Comamonadaceae and Moraxellaceae highlights the importance of monitoring the microbiome. Some members of these families have been reported as opportunistic pathogens.

However, it is essential to note that they also include species with potential applications in biotechnology due to their advantages [64]. The Enterobacteriaceae family comprises members commonly found in the intestines of animals and humans. Fluctuations in the abundance of this family across different months may be linked to variations in the volume of wastewater discharges from homes and hospitals [65]. Finally, the Shewanellaceae family predominantly appeared exclusively at point R4. Considering the characteristics of this family, its presence at the wastewater point is expected, as some of its members are typically found in aquatic environments and play a fundamental role in the decomposition of organic matter [66]. This information emphasizes the diverse array of wastewater bacteria, including pathogenic and beneficial organisms. Furthermore, it emphasizes the concern that discharging wastewater into the Pasto River might be causing a detrimental impact on the environment.

Additionally, it is crucial to highlight species of particular interest, particularly those that may represent a risk to public health due to their pathogenic (Fig. 2A). Our research identified *Bacteroides cellulosilyticus*, widely distributed in the human intestine [67], suggesting fecal contamination in the water. Furthermore, *Raoultella ornithinolytica*, known to be found in bodies of water and even in drinking water systems, as demonstrated by Zou et al. [68], raises concerns as this species has been associated with hospital-acquired infections and in immunosuppressed individuals [68]. Equally important to note is *Pseudomonas aeruginosa*, a species that is not only found in a wide range of environments, including hospitals and sewage systems but also represents a significant public health concern as a nosocomial pathogen and a multi-resistant to antibiotics [69]. The most abundant species identified was *Aeromonas media*, a bacterium capable of causing various diseases in both humans and animals, and it has also been reported to exhibit resistance to specific antibiotics [70]. Lastly, we identified the *Burkholderia cepacia* complex, a group of bacteria widely distributed and highly adaptable, with a tendency for rapid genetic and phenotypic mutations. Despite their environmental origin, these species are of concern as opportunistic pathogens [71]. The identification of these species provides us with valuable tools for regulating the use of water resources and estimating potential risks associated with their presence. It also emphasizes the critical importance of continuously monitoring this resource, as this surveillance is essential for ensuring the control and prevention of public health emergencies. Water contaminated with specific microorganisms can serve as a means of propagating these pathogens, potentially leading to severe illnesses.

Additionally, we would like to highlight species with notable beneficial potential (Fig. 2B), such as *Thauera sp.* *MZIT* is distinguished by its unique ability to produce a specific exopolysaccharide, consequently improving the

viscosity of its environment. This distinctive feature significantly improves the efficiency of disinfection processes in industrial wastewater treatment plants [72]. In contrast, we have also identified *Alicyclophilus denitrificans*, capable of biodegrading a wide range of organic contaminants, suggesting its potential to remediate xenobiotic substances, particularly in contaminated water bodies [73]. Furthermore, among the most abundant species, we have identified *Polaromonas naphthalenivorans*, a bacteria reported in various water bodies that can decompose naphthalene, a common component in fuels, pesticides, and other substances [74, 75]. Our findings go beyond simply identifying species with promising attributes for applications in biotechnology, including bioremediation and the decomposition of harmful compounds. They also shed light on species with potential biotechnology applications and provide valuable information on contaminants that may be present in both wastewater and surface water. This underlines the critical importance of recognizing and characterizing microbial species with substantial potential in these environments.

Additionally, we identified archaea in surface waters, including *Methanobrevibacter smithii*, which is prevalent in the human intestine [76], and *Methanococcus marisaludis*, a marine methanogenic organism [77]. In wastewater, we found *Methanosarcina barkeri*, which has been widely studied as a model of methanogenesis [78]. Regarding viruses, we identified the *Aeromonas virus phiO18P* viruses, known for their ability to transmit resistance and virulence [79], and the *Acanthamoeba polyphaga mimivirus*, responsible for infecting amoebas [80]. Our research highlights a diverse spectrum of microorganisms in wastewater and surface water, some associated with human and animal fecal matter, indicating contamination. We recommend continuing such analyses and improving public databases to collect more complete information on all microorganisms, including parasites and fungi. It would also be highly beneficial to increase the sequencing capacity by including more gigabytes (Gb) of data. This would provide a deeper understanding of the eukaryotic communities, as most of the data is currently allocated to bacterial communities due to their significant diversity.

The generation of metagenome-assembled genomes—MAGs, played a crucial role in identifying previously unknown microorganisms. In environmental samples, MAGs provide essential information about the environment, specific pressures, and the potential to understand non-cultivable or challenging-to-isolate species [89]. This study identified a diverse range of bacterial species, including some originating from the intestine, gastrointestinal system, and feces. These species include *Phocaeicola vulgatus* [90], *Megamonas funiformis* [91], *Prevotella copri* [92], *Bacteroides uniformis* [90], *Phascolarctobacterium_A_succinatutens* [93], *Alistipes putredinis* [94], *Faecalibacterium*

prausnitzii [95] y *Aliarcobacter cryaerophilus_A* [96]. Additionally, species commonly found in water bodies, biofilms, and related environments were detected, such as *Acinetobacter johnsonii* [97], *Sphaerotilus montanus* [98], *Aromonas media* [99], *Aeromonas rivipollensis* [100], *Kaistella chaponensis* [101], *Trichococcus flocculiformis* [102] y *Zoogloea ramigera* [103]. Finally, *Lactococcus A_raffinolactis* is distributed in fermented foods [104]. These findings illustrate the diverse range of genomes within the assembled bacterial species and their respective origins. They contribute to a global understanding that the dominant species originate from the digestive system and those commonly found in the environment.

A key focus of our research centers on the genome of *A. johnsonii*, a microorganism widely distributed in aquatic environments yet also reported as an opportunistic pathogen. Additionally, we note that the category "Information storage and processing" predominates when examining its functional characteristics. This indicates that this species primarily allocates its coding potential to J (translation, ribosomal structure, and biogenesis) and L (replication, recombination, and repair) [47]. Furthermore, assembling the genomes of *Megamonas funiformis* and *Phocaeicola vulgatus* provides a valuable tool to enhance our comprehension of their functions in the human intestine [90, 91]. Regarding functional characteristics (Fig. 3), *P. vulgatus* displays a higher frequency of COGs related to "cell processing and signaling," particularly in M (cell wall/membrane/envelope biogenesis). This underscores the crucial role of this process in maintaining and protecting these species within their environment. *M. funiformis* directs its coding potential toward "metabolism." All of this contributes to a better understanding of the sets of genes that could potentially impact the adaptation and survival of these species in aquatic environments [105, 106]. At a broader level, the most frequent categories were "Unknown function" and "Unclassified," underscoring the importance of exploring and contributing information to public databases to obtain more profound knowledge about these functional attributes. Finally, it is essential to note that genes associated with antibiotic resistance pathways have been identified in all three species, including beta-lactam resistance, vancomycin resistance, and cationic antimicrobial peptide (CAMP) resistance [47]. Therefore, we encourage further comprehensive studies on these functional characteristics, with a specific focus on identifying the precise functions of each gene within these pathways. This effort will allow us to provide comprehensive information on potential risks and shed light on the acquisition of resistance by certain pathogenic species that currently lack detailed characterization (Figs. S4, 3).

Comparative genomics analysis provides valuable insights into the relationships among the samples (Fig. 4). When we conducted the pangenome analysis for the species

A. johnsonii, we observed that reference genome 6 is not clustered with the other reference genomes and the assembled genomes from the study. This could be attributed to the percentage of the core genome identified in this study, which amounts to 218 genes, while the accessory genome accounts for 12.958 genes out of a total of 13.176 genes. This hypothesis is proposed because the reference genome was isolated from water in 2019 [110], and its extensive array of accessory genes might have led to its distinct separation from the others. Furthermore, this species is characterized by having high plasticity in the genome [97].

On the contrary, *M. funiformis* displayed a cluster of diverse samples with no significant grouping differences. However, it is worth noting that this species has a limited number of reference genomes. Lastly, in the case of *P. vulgatus*, sample DP46 from the study appeared distinctly separated from the other samples. This sample was collected from point R4 in wastewater. Nevertheless, it is essential to acknowledge that this genome might be highly fragmented, and its assignment may only partially be accurate. Besides, it may also be influenced by selection pressures such as antibiotic resistance. Likewise, it is essential to highlight the importance of understanding this species since, although it is part of the human intestine, it has stood out for being an opportunistic pathogen [111].

Rivers have encountered significant challenges due to human activities [82, 83], and one of these challenges is pollution caused by the improper disposal of antibiotics. Antibiotics are considered emerging contaminants because they cannot be broken down, leading to their accumulation in the aquatic environment and the development of selection pressure [84, 85]. We examined molecular markers associated with antibiotic resistance (Fig. 5), identifying a high relative abundance related to tetracyclines and aminoglycosides in wastewater. This pattern is consistent with the global context, as both types of drugs are widely used in human, animal, and agricultural healthcare [84]. Furthermore, we noticed a substantial frequency of genes conferring resistance to macrolides, likely linked to their increasing use in livestock and agriculture. As a result, the aquatic environment has become a conduit for the spread of antibiotic-resistant genes [86]. Conversely, in the surface waters of the Pasto River, we identified a high prevalence of genes associated with aminoglycoside and tetracycline resistance, which underscores the potential impact of wastewater discharge directly into surface waters. At point S1, we frequently detected sulfonamide resistance genes, suggesting a notable contribution to the aquatic environment from contaminants such as urine and feces from humans and animals due to their use in medical treatment [87].

In the other analysis, we identified specific molecular markers in wastewater and surface water (Fig. 4). One of these markers is *EmrK*, responsible for conferring resistance

to tetracyclines [81], and *EmrY*, which confers resistance to macrolides, streptogramins, and lincosamides. *Emr* markers are typically found on plasmids, facilitating horizontal gene transfer and enhancing the transmission of these genes to other pathogens [35, 82]. This finding aligns with our previous results, where we observed a high prevalence of genes that confer resistance to tetracyclines. In the case of surface waters, we observed a higher frequency of the *sul1* and *sul2* markers, which are responsible for conferring resistance to sulfonamides. This corresponds with our previous findings, where we identified the gene associated with sulfonamide resistance as commonly present in surface waters [35]. Notably, both markers have been reported in bacterial genera, such as *Acinetobacter*, *Aeromonas*, and *Pseudomonas*, which were also abundant in our study.

Furthermore, identifying molecular markers associated with virulence factors allows us to understand the enzymes, toxins, and other elements that enable pathogenic microorganisms to cause more harm [83]. For instance, we detected markers like *hlyA* [84] in wastewater, which can damage the cell membrane through exotoxin. We also found markers like *mshE* [85], which modify the pilus to enhance adhesion for colonization in aquatic environments and biofilm formation. Additionally, we identified the flagellar transcriptional activator gene *flrC* [86]. In surface waters, we found markers that regulate virulence factors related to stress tolerance, pathogen motility, and more, such as *gacS* [87]. We also identified markers like *sefC* [88], which facilitate the adhesion and colonization of various substrates and cells. While identifying microorganisms in water bodies is crucial, detecting the molecular markers that enhance their virulence, and potential harm is equally essential.

Identifying antibiotic resistance molecular markers from predominant species (Fig. 6), particularly those conferring resistance to aminoglycosides and tetracyclines, raises concerns about the potential public health risks associated with some species, mainly because they are classified as opportunistic pathogens. Moreover, the detection of genes related to resistance to cephamycins, a drug used in the treatment of infections [107], in addition to cephalosporins (which constitute approximately 50% of the antibiotics used in human health) in wastewater, is a cause for concern. This implies the creation of selective pressure in this environment and suggests fecal contamination in surface waters [108]. These findings highlight the inadequate elimination and accumulation of these drugs in water [109]. Regarding markers associated with virulence factors, they were only identified in the genome of *Zoogloea ramigera*. Within this genome, we identified genes such as *hsiCI/vipB* and *hsiCI/vipA*, responsible for tubule-forming proteins of the type VI secretion system *VipB* and type VI *VipA*. *PilT* is associated with the sporadic motility protein *PilT* [36]. In conclusion, while these analyses

provide an initial perspective on markers in the aquatic environment, conducting more comprehensive studies in future research is advisable to obtain robust information for resource monitoring and surveillance.

Conclusion

This study provides vital information about the current state of the Pasto River. We comprehensively analyze the river and its wastewater, explicitly focusing on evaluating physicochemical and microbiological parameters. In essence, this pioneering study is of great importance to the region as it emphasizes the importance of continuous monitoring and surveillance of this critical resource, especially in the context of possible water reuse. Furthermore, this study represents the first microbiological characterization at a regional level, covering both taxonomic and functional aspects, made possible thanks to next-generation sequencing techniques, which provide valuable information on its relevance to public health. This contributes to informed decision-making by relevant authorities responsible for managing these water resources. However, we recommend that future studies incorporate complementary techniques to evaluate the viability of these microorganisms circulating in these bodies of water using next-generation sequencing techniques. Furthermore, paying attention to traditional physicochemical and microbiological analyses is essential to correlate these results with sequencing data. Finally, we encourage the scientific community to continue with the characterization studies of water bodies, and thus nourish public databases to identify and characterize even more microorganisms in future studies.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00284-024-04019-7>.

Author Contribution Vanessa Urrea: Writing—original draft, Conceptualization, Formal analysis, Visualization, Methodology, Data curation, Investigation. Luisa Páez-Triana: Formal analysis, Methodology, Data curation, Investigation. Natalia Velásquez-Ortiz: Writing—review & editing, Methodology, Investigation. Milena Camargo: Writing—review & editing, Methodology, Investigation. Luz H Patiño: Writing—review & editing, Methodology, Investigation. Laura Vega: Visualization, Formal analysis, Data curation. Nathalia Ballesteros: Writing—review & editing, Methodology, Investigation. Arsenio Hidalgo-Troya: Writing—review & editing. Luis-Alejandro Galeano: Writing—review & editing. Juan David Ramírez: Writing—review & editing, Conceptualization, Supervision, Investigation, Funding acquisition. Marina Muñoz: Writing—review & editing, Conceptualization, Supervision, Investigation, Funding acquisition.

Funding Open Access funding provided by Colombia Consortium. This work was financed by the Colombian Fund for Science, Technology, and Innovation of *Sistema General de Regalías* [project

BPIN: 2020000100770]. Neither of the funding entities participated in the study's development.

Data Availability The data obtained and analyzed in this study are available in the European Nucleotide Archive (ENA) repository under project number PRJEB64340.

Declarations

Conflict of Interest The authors declare no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Rajasulochana P, Preethy V (2016) Comparison on efficiency of various techniques in treatment of waste and sewage water—a comprehensive review. *Resour-Effic Technol* 2(4):175–184
2. Kanakoudis V, Tsitsifli S (2017) Potable water security assessment—a review on monitoring, modelling and optimization techniques, applied to water distribution networks. *Desalin Water Treat* 99:18–26. <https://doi.org/10.5004/dwt.2017.21784>
3. Thomson JC (1889) Surface waters. *Am L Rev* 23:372
4. Gad M, Saleh AH, Hussein H, et al (2022) Appraisal of surface water quality of Nile River using water quality indices, spectral signature and multivariate modeling. *Water (Basel)*. <https://doi.org/10.3390/w14071131>
5. Páll E, Niculae M, Kiss T, Şandru CD, Spînu M (2013) Human impact on the microbiological water quality of the rivers. *J Med Microbiol* 62:1635–1640
6. Xu H, Gao Q, Yuan B (2022) Analysis and identification of pollution sources of comprehensive river water quality: evidence from two river basins in China. *Ecol Indic* 135:108561. <https://doi.org/10.1016/j.ecolind.2022.108561>
7. Morin-Crini N, Lichtfouse E, Liu G, Balaram V, Ribeiro ARL, Lu Z, et al. (2022) Worldwide cases of water pollution by emerging contaminants: a review. In: *Environmental chemistry letters*, vol 20. Springer Science and Business Media Deutschland GmbH, pp 2311–2338
8. Lugli GA, Longhi G, Mancabelli L (2022) Tap water as a natural vehicle for microorganisms shaping the human gut microbiome. *Environ Microbiol* 24:3912–3923. <https://doi.org/10.1111/1462-2920.15988>
9. Sobsey MD (2022) Absence of virological and epidemiological evidence that SARS-CoV-2 poses COVID-19 risks from environmental fecal waste, wastewater and water exposures. *J Water Health* 20(1):126–138
10. Kolundžija S, Cheng DQ, Lauro FM (2022) RNA viruses in aquatic ecosystems through the lens of ecological genomics and transcriptomics, vol 14. *Viruses*. MDPI

11. Koderá SM, Sharma A, Martino C, Dsouza M, Grippo M, Lutz L, et al. Microbiome response in an urban river system is dominated by seasonality over wastewater treatment upgrades. <https://doi.org/10.1101/2022.06.30.498375>
12. Amarasiri M, Sano D, Suzuki S (2020) Understanding human health risks caused by antibiotic resistant bacteria (ARB) and antibiotic resistance genes (ARG) in water environments: Current knowledge and questions to be answered. *Crit Rev Environ Sci Technol* 50(19):2016–2059
13. Feng C, Xu Z, Li Y, Zhu N, Wang Z (2021) Research progress on the contamination status and control policy of *Giardia* and *Cryptosporidium* in drinking water. In: *Journal of water sanitation and hygiene for development*, vol 11. IWA Publishing, pp 867–886
14. Zhu K, Ren H, Lu Y (2022) Potential biorisks of *Cryptosporidium* spp. and *Giardia* spp. from reclaimed water and countermeasures. In: *Current pollution reports*. Springer Science and Business Media Deutschland GmbH
15. Rothman JA, Loveless TB, Kapcia J (2021) RNA viromics of Southern California wastewater and detection of SARS-CoV-2 single-nucleotide variants. *Appl Environ Microbiol* 87:e01448. <https://doi.org/10.1128/AEM.01448-21>
16. Song W, Chen H, Xue N, et al (2022) Metagenomic binning and assembled genome analysis revealed the distinct composition of resistome and mobilome in the Lli River. *Ecotoxicol Environ Saf* 242:113886. <https://doi.org/10.1016/j.ecoenv.2022.113886>
17. Payne JT, Millar JJ, Jackson CR, Ochs CA (2017) Patterns of variation in diversity of the Mississippi river microbiome over 1300 kilometers. *PLoS One* 12:1–18. <https://doi.org/10.1371/journal.pone.0174890>
18. Sutherland BJ, Finke JF, Saunders R, Warne S, Schulze AD, Strohm T, et al. Metatranscriptomics reveals a shift in microbial community composition and function during summer months in a coastal marine environment. <https://doi.org/10.1101/2022.01.21.477286>
19. Chopyk J, Nasko DJ, Allard S, et al (2020) Comparative metagenomic analysis of microbial taxonomic and functional variations in untreated surface and reclaimed waters used in irrigation applications. *Water Res* 169:115250. <https://doi.org/10.1016/j.watres.2019.115250>
20. Tang Y, Liang Z, Li G, et al (2021) Metagenomic profiles and health risks of pathogens and antibiotic resistance genes in various industrial wastewaters and the associated receiving surface water. *Chemosphere* 283:131224. <https://doi.org/10.1016/j.chemosphere.2021.131224>
21. Kim JJ, Seong HJ, Johnson TA, et al (2023) Persistence of antibiotic resistance from animal agricultural effluents to surface water revealed by genome-centric metagenomics. *J Hazard Mater* 457:131761. <https://doi.org/10.1016/j.jhazmat.2023.131761>
22. Scherer GS, Guiguet Leal DA, Goulart JAG, Araújo RS, Beux MR, Moreira NM (2022) Parasitological, microbiological, and antimicrobial resistance profiles of raw and drinking water in a tourist city in the tri-border region of South America. *J Water Health* 20(2):385–395
23. Breton-Deval L, Juárez K, Sánchez-Flores A, Vera-Estrella R. Water quality assessments and metagenomic analysis of the polluted river Apatlaco, Mexico. <https://doi.org/10.1101/536128>
24. Rodríguez EA, Ramirez D, Balcázar JL, Jiménez JN (2021) Metagenomic analysis of urban wastewater resistome and mobilome: a support for antimicrobial resistance surveillance in an endemic country. *Environ Pollut* 276:116736. <https://doi.org/10.1016/j.envpol.2021.116736>
25. Lucía M, Martínez O, Carolina A, et al (2022) Instituto Nacional De Salud. <https://www.ins.gov.co/BibliotecaDigital/Boletin-vigilancia-calidad-agua-mayo-2022.pdf>. Accessed 25 Sep 2023
26. Sánchez C, López MC, Galeano LA, et al (2018) Molecular detection and genotyping of pathogenic protozoan parasites in raw and treated water samples from southwest Colombia. *Parasit Vectors* 11:563. <https://doi.org/10.1186/s13071-018-3147-3>
27. Empopasto (2021) Cuentas de abastecimiento. Cuencas de abastecimiento. <https://emposig.empopasto.com.co/arcgisportal/apps/experiencebuilder/experience/?id=d36adf0ed3b147daaa22d9be9f874131&page=Home&views=Cueca-Pasto>. Accessed 11 Jul 2024
28. Corponariño (2011) Programa manejo integral del recurso hídrico en el departamento de nariño. Plan de Ordenación del Río Pasto
29. Lauritsen JM, Bruus ME (2003) A comprehensive tool for validated entry and documentation of data. The EpiData Association, Odense, Denmark, 2005
30. Andrews S (2010) <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>. FastQC: A Quality Control Tool for High Throughput Sequence Data
31. Ewels P, Magnusson M, Lundin S, Käller M (2016) MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* 32(19):3047–3048
32. Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30(15):2114–2120
33. Kim D, Song L, Breitwieser FP, Salzberg SL (2016) Centrifuge: rapid and sensitive classification of metagenomic sequences. *Genome Res* 26(12):1721–1729
34. Breitwieser FP, Salzberg SL (2020) Pavian: interactive analysis of metagenomics data for microbiome studies and pathogen identification. *Bioinformatics* 36(4):1303–1304
35. Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A et al (2020) CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucl Acids Res* 48(D1):D517–D525
36. Liu B, Zheng D, Zhou S, Chen L, Yang J (2022) VFDB 2022: a general classification scheme for bacterial virulence factors. *Nucl Acids Res* 50(D1):D912–D917
37. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA (2017) MetaSPAdes: a new versatile metagenomic assembler. *Genom Res* 27(5):824–834
38. Kang DD, Froula J, Egan R, Wang Z (2015) MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ* 3:e1165. <https://doi.org/10.7717/peerj.1165>
39. Wu YW, Simmons BA, Singer SW (2016) MaxBin 20: an automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 32(4):605–607
40. Alneberg J, Bjarnason BS, de Bruijn I, et al (2014) Binning metagenomic contigs by coverage and composition. *Nat Methods* 11:1144–1146. <https://doi.org/10.1038/nmeth.3103>
41. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG et al (2018) Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat Microbiol* 3(7):836–843
42. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW (2015) CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25(7):1043–1055
43. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH (2020) GTDB-Tk: a toolkit to classify genomes with the genome taxonomy database. *Bioinformatics* 36(6):1925–1927
44. Seemann T (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30(14):2068–2069
45. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG et al (2015) Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* 31(22):3691–3693

46. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J (2021) eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol Biol Evol* 38(12):5825–5829
47. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S et al (2020) KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. *Bioinformatics* 36(7):2251–2252
48. Sammarro Silva KJ, Sabogal-Paz LP (2021) *Cryptosporidium* spp. and *Giardia* spp. (oo)cysts as target-organisms in sanitation and environmental monitoring: a review in microscopy-based viability assays. In: *Water research*, vol 189. Elsevier
49. Wood DE, Lu J, Langmead B (2019) Improved metagenomic analysis with Kraken 2. *Genome Biol* 20(1)
50. Zhu L, Li R, Yan Y, Cui L (2022) Urbanization drives the succession of antibiotic resistome and microbiome in a river watershed. *Chemosphere* 301:134707. <https://doi.org/10.1016/j.chemosphere.2022.134707>
51. Ustaoglu F, Tepe Y, Taş B (2020) Assessment of stream quality and health risk in a subtropical Turkey river system: a combined approach using statistical analysis and water quality index. *Ecol Indic* 113:105815. <https://doi.org/10.1016/j.ecolind.2019.105815>
52. Verma A (2014) Assessment of water quality and Pollution Status of Kalpi (Morar) River, Gwalior, Madhya Pradesh: with special reference to Conservation and Management plan Water Treatment and Water Technology View project [Internet]. <https://www.researchgate.net/publication/266065376>
53. Pattanayak S, Priyadarsini S, Paul A, et al (2020) Diversity of virulence-associated genes in pathogenic *Aeromonas hydrophila* isolates and their in vivo modulation at varied water temperatures. *Microb Pathog* 147:104424. <https://doi.org/10.1016/j.micpath.2020.104424>
54. Yu Q, Han Q, Shi S (2023) Metagenomics reveals the response of antibiotic resistance genes to elevated temperature in the Yellow River. *Sci Total Environ* 859:160324. <https://doi.org/10.1016/j.scitotenv.2022.160324>
55. Ramo A, Del Cacho E, Sánchez-Acedo C, Quílez J (2017) Occurrence and genetic diversity of *Cryptosporidium* and *Giardia* in urban wastewater treatment plants in north-eastern Spain. *Sci Total Environ* 598:628–638
56. Lass A, Szostakowska B, Korzeniewski K, Karanis P (2017) Detection of *Giardia intestinalis* in water samples collected from natural water reservoirs and wells in northern and north-eastern Poland using LAMP, real-time PCR and nested PCR. *J Water Health* 15(5):775–787
57. Hemati S, Mohammadi-Moghadam F, Mohammadian-Hafshejani A, et al (2022) Occurrence of *Giardia* and *Cryptosporidium* in effluents of urban wastewater treatment plants: a global systematic review and meta-analysis. *J Clean Prod* 378:134555. <https://doi.org/10.1016/j.jclepro.2022.134555>
58. Zhang L, Ben Said L, Hervé N, et al (2022) Effects of drinking water supplementation with *Lactobacillus reuteri*, and a mixture of reuterin and microcin J25 on the growth performance, caecal microbiota and selected metabolites of broiler chickens. *J Anim Sci Biotechnol*. <https://doi.org/10.1186/s40104-022-00683-6>
59. Jean S, Wallace MJ, Dantas G (2022) Burnham CA (2022) Time for some group therapy: update on identification, antimicrobial resistance, taxonomy, and clinical significance of the *Bacteroides fragilis* group. *J Clin Microbiol* 60(9):e02361-20
60. Chernitsyna S, Elovskaya I, Pogodaeva T, et al (2023) Bacterial communities in a gradient of abiotic factors near a sulfide thermal spring in Northern Baikal. *Diversity (Basel)*. <https://doi.org/10.3390/d15020298>
61. Kerber-Diaz JC, Leos-Ramírez MA, Flores-Ceron AA, et al (2022) Distribution of CRISPR-Cas systems in the Burkholderiaceae family and its biological implications. *Arch Microbiol* 204:703. <https://doi.org/10.1007/s00203-022-03312-y>
62. Wang Z, Li W, Li H, Zheng W, Guo F (2022) Phylogenomics of Rhodocyclales and its distribution in wastewater treatment systems. *Sci Rep* 10(1)
63. Amin SQ, Mahmood HJ, Zorab HK (2023) *Campylobacteriosis*. In: *One health triad*, vol 2. Unique Scientific Publishers, Faisalabad, Pakistan [Internet], pp 87–93. <https://doi.org/10.47278/book.oht/2023.46>
64. Knoot CJ, Wantuch PL, Robinson LS, Rosen DA, Scott NE, Harding CM (2023) Discovery and characterization of a new class of O-linking oligosaccharyltransferases from the Moraxellaceae family. *Glycobiology* 33(1):57–74
65. Singh AK, Das S, Kumar S (2020) Distribution of antibiotic-resistant enterobacteriaceae pathogens in potable spring water of eastern indian himalayas: emphasis on virulence gene and antibiotic resistance genes in *Escherichia coli*. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2020.581072>
66. Xu S, Yu K, Su H, Chen B, Huang W, Liang J et al (2019) Proposal of *Parashewanella* gen. nov. to accommodate *Parashewanella curva* sp. nov. and *Parashewanella spongiae* comb. nov. in the Shewanellaceae. *Int J Syst Evol Microbiol* 69(5):1259–1264
67. Christensen L, Sørensen CV, Wøhlk FU, et al (2020) Microbial enterotypes beyond genus level: bacteroides species as a predictive biomarker for weight change upon controlled intervention with arabinoxylan oligosaccharides in overweight subjects. *Gut Microbes*. <https://doi.org/10.1080/19490976.2020.1847627>
68. Zou H, Berglund B, Xu H, et al (2020) Genetic characterization and virulence of a carbapenem-resistant *Raoultella ornithinolytica* isolated from well water carrying a novel megaplasmid containing bla_{NDM-1}. *Environ Pollut* 260:114041. <https://doi.org/10.1016/j.envpol.2020.114041>
69. Tuon FF, Dantas LR, Suss PH, Tasca Ribeiro VS (2022) Pathogenesis of the *Pseudomonas aeruginosa* biofilm: a review. *Pathogens* 11
70. Talagrand-Reboul E, Roger F, Kimper JL, et al (2017) Delineation of taxonomic species within complex of species: aeromonas media and related species as a test case. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2017.00621>
71. Tavares M, Kozak M, Balola A, Sá-Correia I (2020) *Burkholderia cepacia* complex bacteria: a feared contamination risk in water-based pharmaceutical products. <https://doi.org/10.1128/CMR>
72. Jiang K, Sanseverino J, Chauhan A, Lucas S, Copeland A, Lapidus A et al (2012) Complete genome sequence of *Thauera aminoaromatica* strain MZ1T. *Stand Genom Sci* 6(3):325–335
73. Fuentes-Jaime J, Vargas-Suárez M, Cruz-Gómez MJ, Loza-Tavera H (2022) Concerted action of extracellular and cytoplasmic esterase and urethane-cleaving activities during Impranol biodegradation by *Alicyclophilus denitrificans* BQ1. *Biodegradation* 33(4):389–406
74. Padilla-Del Valle R, Morales-Vale LR, Ríos-Velázquez C (2017) Unraveling the microbial and functional diversity of Coamo thermal spring in Puerto Rico using metagenomic library generation and shotgun sequencing. *Genom Data* 11:98–101
75. Hwang K, Choe H, Nasir A, Kim KM (2021) Complete genome of *Polaromonas vacuolata* KCTC 22033T isolated from beneath Antarctic Sea ice. *Mar Genom* 55:100790. <https://doi.org/10.1016/j.margen.2020.100790>
76. Bang C, Weidenbach K, Gutschmann T (2014) The intestinal archaea methanosphaera stadmanae and methanobrevibacter smithii activate human dendritic cells. *PLoS One* 9:e99411
77. Palabikyan H, Ruddyard A, Pomper L (2022) Scale-up of biomass production by *Methanococcus maripaludis*. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2022.1031131>

78. Ye J, Yu J, Zhang Y (2019) Light-driven carbon dioxide reduction to methane by *Methanosarcina barkeri*-CdS biohybrid. *Appl Catal B* 257:117916 <https://doi.org/10.1016/j.apcatb.2019.117916>
79. Subirats J, Sánchez-Melsió A, Borrego CM, Balcázar JL, Simonet P (2016) Metagenomic analysis reveals that bacteriophages are reservoirs of antibiotic resistance genes. *Int J Antimicrob Agents* 48(2):163–167
80. Ferek JD, Thoden JB, Holden HM (2020) Biochemical analysis of a sugar 4,6-dehydratase from *Acanthamoeba polyphaga* Mimivirus. *Protein Sci* 29(5):1148–1159
81. Christaki E, Marcou M, Tofarides A (2020) Antimicrobial resistance in bacteria: mechanisms, evolution, and persistence. In: *Journal of molecular evolution*, vol 88. Springer, pp 26–40
82. Abushaheen MA, Muzaheed Fatani AJ, et al (2020) Antimicrobial resistance, mechanisms and its clinical significance. *Disease-a-Month* 66:100971. <https://doi.org/10.1016/j.disamonth.2020.100971>
83. Leitão JH (2020) Microbial virulence factors. In: *International journal of molecular sciences*, vol 21. MDPI AG, pp 1–6
84. Noori AA (2023) Detection of the, hlyA, fimH PgaD gene in *Escherichia coli* isolated from different sources in Kirkuk city Sundus Jassim Mohamad
85. Zhang YL, Wu F, Huang YC, Jian JC, Cai SH (2023) Mechanisms underlying the virulence and intestinal colonization of the *Vibrio alginolyticus* HY9901 DctP protein with proteomic analysis. *Appl Biochem Microbiol* [Internet] 59(5):646–658. <https://doi.org/10.1134/S0003683823050198>
86. Syed KA, Beyhan S, Correa N, Queen J, Liu J, Peng F et al (2009) The *Vibrio cholerae* flagellar regulatory hierarchy controls expression of virulence factors. *J Bacteriol* 191(21):6555–6570
87. Strateva T, Mitov I (2011) Contribution of an arsenal of virulence factors to pathogenesis of *Pseudomonas aeruginosa* infections. *Ann Microbiol* 61:717–732
88. El-Tayeb MA, Ibrahim ASS, Maany DA, Elbadawi YB, Al-Salamah AA (2019) Detection of virulence genes and investigation of the immunogenicity of outer membrane proteins of salmonella enterica. *J Appl Biol Biotechnol* 7(5):62–73
89. Pinhasi J, Farnelid H, García SM (2022) Functional responses of key marine bacteria to environmental change—toward genetic counselling for coastal waters. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2022.869093>
90. Neff A, Lück R, Hövels M, Deppenmeier U (2023) Expanding the repertoire of counterselection markers for markerless gene deletion in the human gut bacterium *Phocaeicola vulgatus*. *Anaerobe* 1:81
91. Chen G, Zeng Z, Xie M, Peng Y, Zhou W, Xu W et al (2022) Fermentation characteristics and probiotic activity of a purified fraction of polysaccharides from Fuzhuan brick tea. *Food Sci Hum Wellness* 11(3):727–737
92. Abdelsalam NA, Hegazy SM, Aziz RK (2023) The curious case of *Prevotella copri*. In: *Gut Microbes*, vol 15. Taylor and Francis
93. Shigeno Y, Kitahara M, Shime M, Benno Y (2019) *Phascolarctobacterium wakonense* sp. nov., isolated from common marmoset (*Callithrix jacchus*) faeces. *Int J Syst Evol Microbiol* 69(7):1941–1946
94. Tyrrell KL, Warren YA, Citron DM, Goldstein EJC (2011) Reassessment of phenotypic identifications of *Bacteroides putredinis* to *Alistipes* species using molecular methods. *Anaerobe* 17(3):130–134
95. Leylabadlo HE, Ghotaslou R, Feizabadi MM, Farajnia S, Moadab SY, Ganbarov K et al (2020) The critical role of *Faecalibacterium prausnitzii* in human health: an overview. *Microb Pathog* 1:149
96. Gungor C, Hizlisoy H, Ertas Onmaz N, et al (2023) Profile of *Aliarcobacter* spp. from edible giblets: genetic diversity, antibiotic resistance, biofilm formation. *Int J Food Microbiol* 386:110047. <https://doi.org/10.1016/j.ijfoodmicro.2022.110047>
97. Jia J, Liu M, Feng L, Wang Z (2022) Comparative genomic analysis reveals the evolution and environmental adaptation of *Acinetobacter johnsonii*. *Gene* 15:808
98. Nott MA, Driscoll HE, Takeda M, et al (2020) Advanced biofilm analysis in streams receiving organic deicer runoff. *PLoS One* 15:e0227567
99. Dubey S, Ager-Wick E, Peng B, et al (2022) Characterization of virulence and antimicrobial resistance genes of *Aeromonas media* strain SD/21–15 from marine sediments in comparison with other *Aeromonas* spp. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2022.1022639>
100. Fono-Tamo EUK, Kamika I, Dewar JB, Lekota KE (2023) Comparative genomics revealed a potential threat of *Aeromonas rivipollensis* G87 strain and its antibiotic resistance. *Antibiotics*. <https://doi.org/10.3390/antibiotics12010131>
101. Ai J, He X, Ren M, et al (2024) *Kaistella yananensis* sp. Nov., a novel indoleacetic acid-producing bacterium isolated from the root nodules of *Sophora davidii* (Franch.) skeels. *Curr Microbiol* 81:60. <https://doi.org/10.1007/s00284-023-03578-5>
102. Jin-gon S, Shin-Gyu C, Se-Hwan K, et al (2022) Heliorhodopsin helps photolyase to enhance the DNA repair capacity. *Microbiol Spectr* 10:e02215–22. <https://doi.org/10.1128/spectrum.02215-22>
103. Bharti M, Nagar S, Khurana H, Negi RK (2022) Metagenomic insights to understand the role of polluted river Yamuna in shaping the gut microbial communities of two invasive fish species. *Arch Microbiol* 204:509. <https://doi.org/10.1007/s00203-022-03127-x>
104. Jung MY, Lee C, Seo M-J, et al (2020) Characterization of a potential probiotic bacterium *Lactococcus raffinolactis* WiKim0068 isolated from fermented vegetable using genomic and in vitro analyses. *BMC Microbiol* 20:136. <https://doi.org/10.1186/s12866-020-01820-9>
105. Zhang W, Xu X, Cai L, Cai X (2023) Dysbiosis of the gut microbiome in elderly patients with hepatocellular carcinoma. *Sci Rep* 13:7797. <https://doi.org/10.1038/s41598-023-34765-w>
106. Cardos HB, Dalmacio LM, Balolong MP. Bacterial composition and antibiotic resistome profile of water in the Manila Bay South Harbor [Internet]. <https://www.researchgate.net/publication/373238250>
107. Yamashiro H, Kasamatsu Y, Anan N, Takemura M, Yamano Y (2023) In vitro efficacy of humanized regimen of flomoxef against extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 67(9)
108. Das N, Madhavan J, Selvi A, Das D (2019) An overview of cephalosporin antibiotics as emerging contaminants: a serious environmental concern. *3 Biotech*, vol 9. Springer
109. Li J, Li W, Liu K, Guo Y, Ding C, Han J, et al. (2022) Global review of macrolide antibiotics in the aquatic environment: sources, occurrence, fate, ecotoxicity, and risk assessment. In: *Journal of hazardous materials*, vol 439. Elsevier B.V.
110. Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ et al (2023) Introducing the bacterial and viral bioinformatics resource center (BV-BRC): a resource combining PATRIC, IRD and ViPR. *Nucl Acids Res* 51(1D):678–689
111. Vu H, Muto Y, Hayashi M, Noguchi H, Tanaka K, Yamamoto Y (2022) Complete genome sequences of three *Phocaeicola vulgatus* strains isolated from a healthy Japanese individual. *Microbiol Resour Announc* 11(2):e01124-21