

Blastocystis genetic diversity in animal and human samples from different departments of Colombia using complete sequencing of the 18S rRNA gene (SSU rRNA) by Oxford Nanopore Technologies (ONT)

Paula Jiménez^a, Marina Muñoz^a, Lissa Cruz-Saavedra^a, Anny Camargo^{a,b}, Juan David Ramírez^{a,c,*}

^a Centro de Investigaciones en Microbiología y Biotecnología-UR (CIMBIUR), Facultad de Ciencias Naturales, Universidad del Rosario, Bogotá, Colombia

^b Health Sciences Faculty, Universidad de Boyacá, Tunja, Colombia

^c Molecular Microbiology Laboratory, Department of Pathology, Molecular and Cell-based Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, United States

ARTICLE INFO

Keywords:
Blastocystis
 Subtypes
 STs
 Colombia
 Animals
 Humans
 SSU rRNA
 Sequencing

ABSTRACT

Blastocystis is an intestinal microeukaryote that has raised attention due to its wide distribution in animals and humans. The risk of zoonotic circulation primarily arises from close contact with infected animals. Therefore, the following study aimed to evaluate the diversity and frequency of *Blastocystis* subtypes in Colombian human and animal samples using complete sequencing of the 18S rRNA gene. For this purpose, 341 human stool samples and 277 animal fecal samples (from cattle, sheep, goat, pigs, cats, and dogs), were collected from different Colombian regions and analyzed using PCR-based detection and full-length 18S SSU rRNA gene Next-Generation Sequencing (NGS). Among the 618 samples from both hosts, humans and animals, the results revealed widespread *Blastocystis* frequency, with 48.09% ($n = 164$) in humans and 31.4% ($n = 87$) detection in animals. Dogs, cats, sheep, pigs, and wild animals tested positive, aligning with global prevalence patterns. Also, 29 human samples and 23 animal samples were sequenced using ONT technology from which 11 long-read unique sequences were generated and cluster with their compared reference sequences. The subtype distribution varied within hosts, detecting ST1 and ST3 in both human and animal samples. Subtypes ST5, ST10, ST14, ST15, ST21, ST24, ST25 and ST26 were limited to animals hosts, some of which are considered to have zoonotic potential. On the other hand, ST2 was found exclusively in human samples from Bolivar region. Mixed infections occurred in both animal and humans, 60.86% and 27.58% respectively. Moreover, to our knowledge, this is the first study in Colombia identifying ST15 in pigs and ST25 in sheep. The subtypes (STs) identified in this study indicate that certain animals may serve as reservoirs with the potential for zoonotic transmission. The identification of zoonotic subtypes highlights the use of Next Generation Sequencing as the depth and resolution of the sequences increases providing insights into STs of medical and veterinarian significance. It also reveals the coexistence of diverse subtypes among hosts. Further research is essential for understanding transmission dynamics, health implications, and detection strategies for *Blastocystis* occurrence in animals and humans, mainly associated to the role of animals as reservoirs and their close interaction with humans.

1. Introduction

Blastocystis, an enteric protist classified within the chromista kingdom and recognized as a stramenopile (Stensvold et al., 2020), predominantly colonizes the gastrointestinal tract of diverse hosts (Qi et al., 2020; Yoshikawa et al., 2016). It exhibits a widespread global distribution, with prevalence estimates ranging from 0.5% to 24% in

developed nations and escalating to 50–100% in developing countries (Alfellani et al., 2013a). The primary transmission route involves the fecal-oral pathway (Lee et al., 2012; Li et al., 2007), facilitated by water, food, and contact with certain animals as the most probable propagation mechanisms (Londoño-Franco et al., 2014; Maloney and Santin, 2021).

While the manifestation of symptoms due to *Blastocystis* infections in animals remains unclear, a substantial prevalence of subtypes has been

* Corresponding author.

E-mail addresses: juand.ramirez@urosario.edu.co, juan.ramirezgonzalez@mssm.edu (J.D. Ramírez).

<https://doi.org/10.1016/j.actatropica.2023.107090>

Received 18 September 2023; Received in revised form 27 November 2023; Accepted 1 December 2023

Available online 2 December 2023

0001-706X/© 2023 Elsevier B.V. All rights reserved.

documented across various species. Noteworthy, some reports provide evidence for *Blastocystis* potential zoonotic transmission between humans and animals (Hublin et al., 2020; Ramírez et al., 2017). In contrast, among human populations, *Blastocystis* has been identified as a possible cause of gastrointestinal and cutaneous symptoms. Interestingly, its frequent identification in asymptomatic patients adds complexity to understanding the intricate relationship between infection and disease (Ajajampur and Tan, 2016; Casero et al., 2015; Clark et al., 2013; Wawrzyniak et al., 2013). Additionally, there are ongoing investigations into whether it acts as a pathogen or as a part of the healthy intestinal microbiota (Deng et al., 2021).

This microeukaryote exhibits a wide genetic diversity (C. Rune Stensvold et al., 2012; Stensvold et al., 2020), currently 40 subtypes (ST) have been reported, based on nucleotide polymorphism within the small subunit of ribosomal RNA (rRNA-18S) (Hernández-Castro et al., 2023; Higuera et al., 2021; Rondón et al., 2023; Santin et al., n.d.; Stensvold, 2013; Stensvold et al., 2023; Stensvold and Clark, 2020). Certain subtypes display moderate specificity towards human and animal hosts, while others are exclusively identified in animals (Alfellani et al., 2013a; Barbosa et al., 2017; Ramirez et al., 2016). ST1 to ST4, are predominantly detected from humans (90%). Meanwhile, ST5 to ST8 are primarily associated with animal hosts, such as sheep, pigs, dogs, cats, rodents, and birds. Remarkably, ST5 has also been detected in amphibians and was recently reported in human samples within a Colombian population by Vega et al. (Alfellani et al., 2013b; Vega Romero, 2020). Conversely, ST9 has exclusively reported in human hosts (Stensvold and Clark, 2016). ST10 to ST17 predominantly inhabit non-human hosts, including marsupials, goats, non-human primates, elephants, wild birds, and sheep (Maloney et al., 2018, 2021a; Roberts et al., 2013). Although subtypes ST12, ST13, and ST16, are commonly associated with non-human hosts, researchers such as Ramírez et al. and Osorio-Pulgarín et al. have detected them in human samples (P.A. Jiménez et al., 2019; Osorio-Pulgarín et al., 2021; Ramirez et al., 2016). ST21 to ST26 have been documented in cattle and sheep, ST21 was identified in antelopes (*Kobus ellipsiprymnus*) (Maloney et al., 2021a; Zhao et al., 2017) and ST24 in Llamas (*Lama glama*) (Higuera et al., 2021). ST27 and ST28 were documented in samples from wild birds (Maloney et al., 2020a), while ST29 was detected in chickens (Maloney et al., 2021a). Recent additions include ST30 and ST31, characterized in white-tailed deer (*Odocoileus virginianus*) (Maloney et al., 2021b). Additionally, Higuera et al. reported subtype ST32 in Colombian cattle and goats (Higuera et al., 2021), whereas ST33 and ST34 were found in Colombian horses (Baek et al., 2022). Novel ST41 identified in a Colombian patient undergoing a colorectal cancer screening (Hernández-Castro et al., 2023). Furthermore, genetic diversity varies considerably among subtypes and even within individual subtypes, reflecting pronounced genetic heterogeneity which could be associated with biological features and phenotypic effect on the host (Stensvold, 2013).

Molecular characterization of *Blastocystis* involves techniques such as PCR-based partial amplification of the rRNA-18S gene (600 pb), followed by Sanger sequencing, which leads to subtype assignment via database comparison. Nonetheless, this method is known for missing mixed infections by omitting subtypes in low abundance, leading to underestimated results in terms of subtype diversity. Conversely, Illumina sequencing has been adopted for *Blastocystis* subtyping, demonstrating heightened sensitivity and specificity, especially in scenarios involving mixed infections, low abundance. This technique notably provides valuable information of those subtypes with zoonotic potential such as ST1 to ST5 (Maloney et al., 2019).

Emerging third-generation sequencing methods, such as Oxford Nanopore Technologies (ONT) using the MinION device, have been used to obtain complete sequences of the *Blastocystis* 18S SSU rRNA gene (1800 pb), increasing the accuracy of genetic information obtained, enhancing confidence in the identified subtypes. The MinION's unique attributes, like generating long reads of up to 882 kb and avoiding the use of multiple primers or repeated PCRs, enabling the identification of

novel subtypes and proper phylogenetic analysis (Jain et al., 2018; Maloney and Santin, 2021). Furthermore, MinION reduces sequencing duration from days to hours, diminishes PCR-induced biases, and continuous improvements sequence processing and software lead to accurate individual reads up to 99% (Bowden et al., 2019; Hernández et al., 2023; Katoh and Standley, 2013; Maloney et al., 2020b).

The generation of full-length 18S SSU rRNA gene sequences not only offers reference material for subtype identification and robust phylogenetic lineage development but also proves valuable for designating new subtypes. The complete gene sequence meets the 4% minimum divergence threshold, preventing invalid subtype creation or nomenclatural errors (Higuera et al., 2021; Stensvold, 2013; Stensvold and Clark, 2020). Indeed, during the last years with this technology it was possible to identify eight new subtypes, such as ST32 in Colombian goats and cows, ST33 and ST34 in Colombian horses (Baek et al., 2022), among others up to ST38 (Hernández et al., 2023; Maloney et al., 2023). This methodology facilitates a deeper understanding of mixed infections within single samples, reducing the necessity of obtaining individual reference sequences via pure cultures for each subtype. Its utility improve the understanding of transmission dynamics, pathogenicity, and host specificity in a genetically diverse microorganism like *Blastocystis* (Maloney et al., 2020b).

In Colombia, few studies have focused on the molecular characterization of *Blastocystis* within animal and human populations using ONT. While the National Survey of Intestinal Parasitology suggests a prevalence range of 3.6% to 60.3% (Ministerio de Salud y Protección Social, 2015), distinct distributions of subtypes have been identified across departments such as Guainía, Quindío, Tolima, Cundinamarca, Boyacá, Amazonas, Cauca, Antioquia, Bolívar, Córdoba and Casanare (Castañeda et al., 2020; Higuera et al., 2020a; P.A. Jiménez et al., 2019; Ramirez et al., 2016; Vega Romero, 2020; Villamizar et al., 2019). Predominantly, Andean, and Caribbean regions exhibit high diversity and prevalence of intestinal parasites, including *Blastocystis* (Higuera et al., 2020b). However, research has primarily focused on intestinal parasitism linked to nutritional deficiencies and infantile polyparasitism (Osorio-Pulgarín et al., 2021; Ramirez et al., 2017).

The molecular characterization of intestinal protists such as *Blastocystis* is essential for understanding its epidemiology, and frequency, comprehending transmission dynamics and host specificity. Due to its high frequency, *Blastocystis* play a significant role in public health and disease research implications. Despite the elevated *Blastocystis* prevalence in Colombia, most studies have employed PCR and sequencing techniques (Sanger or Illumina) to analyze a fragment of the SSU rRNA gene, resulting in an increasing focus on animal hosts and a decreasing emphasis on human subtyping. However, comprehensive gene sequences remain sparse, with limited studies exploring both host types. Oxford Nanopore's MinION technology has bridged this gap in animal hosts, even though with limited human research or combined host analyses. Consequently, complete rRNA-18S gene sequences of diverse *Blastocystis* subtypes in Colombia remain unavailable. Therefore, this study seeks to address this gap by characterizing the genetic and epidemiological diversity of *Blastocystis* in both animal and human hosts across diverse Colombian regions, thereby generating full-length rRNA-18S gene sequences.

2. Materials and methods

2.1. Study population and sample collection

The human samples were originally collected in a previous study during which the identification of *Blastocystis* was also carried out, but were utilized in the current study. A total of 341 stool samples were collected from individuals ranging various age groups from 1 to 70 years old, (mean: 5 years - standard deviation: 6 years) residing in both rural and urban settings. Convenience sampling was conducted across diverse regions of Colombia, including departments such as Casanare ($n = 53$),

Cauca ($n = 258$), and Bolívar ($n = 30$) (Higuera et al., 2020b).

Additionally, fecal samples were included from 277 apparently healthy domestic animals and livestock representing distinct species. These samples were collected from rural areas in Boyacá (Jenesano, Gámeza and Paipa) and urban areas in Bogotá, including Pigs (*Sus scrofa*) ($n = 36$), Cattle (*Bos taurus*) ($n = 29$), Sheep (*Ovis aries*) ($n = 38$), Goats (*Capra aegagrus hircus*) ($n = 20$), Cats (*Felis catus*) ($n = 96$) Dogs (*Canis lupus familiaris*) ($n = 35$), these animals were apparently healthy and have no exhibit digestive symptoms. Sampling was conducted during the period from November to March 2022, large animal (cows, pigs, sheep, and goats) sampling was performed by anal conduct collection, around 300 mg of stool were collected in transparent sterile container for coprological analysis. In the case of small animals such as cats and dogs with minimal contact between the animal and the keepers, around 200 mg of stool was collected from the ground using transparent sterile container for coprological analysis.

Also, stool samples were collected from zoo animals ($n = 23$) whose species are shown in Supplementary Table 2. These samples were collected in November 2019 from wild mammals rescued from illegal trafficking and held in captivity in Cubará, Boyacá. Despite their previous exposure to humans and other animals, these animals exhibited no signs of intestinal infection. They were housed in dedicated enclosures and provided with a specific diet under veterinary supervision and had access to cistern water sourced from a stream-connected reservoir. No deworming protocol had been initiated previously. Stool samples were collected both individually and in groups (from animals of the same species housed together detailed in Supplementary Table 2), with minimal contact between the animals and keepers, approximately 200 mg of stool were collected from the ground using Eppendorf tubes containing ethanol (Cruz-Saavedra et al., 2023).

The specific geographical origins of these samples are displayed in Fig. 1. All samples were stored at $-30\text{ }^{\circ}\text{C}$ until DNA extraction without any preservative intervention. Samples that were unavailable during molecular analysis were excluded, and due to the absence of accompanying clinical information, clinical parameters were not established as exclusion criteria.




Fig. 1. Geographic Distribution of Sample Collection. Exact geographical location from the municipalities where the samples were collected are indicated in dots of distinct colors by department. The total number of samples and host from which those were taken are exhibited in the Location legend.

2.2. DNA extraction and Blastocystis detection via PCR

DNA extraction and *Blastocystis* detection had been previously performed in human samples by Higuera et al., resulting in the identification of 164 positive samples (Higuera et al., 2020b). Subsequently, a survey was conducted to assess the current availability of these samples, revealing that 111 of them were still available (Supplementary Table 1). For animal samples, prior to DNA extraction, 300 μL of the stool were washed with sterile saline and phosphate buffer (PBS 1X) three times. Subsequently, 600 mL of PBS was added to each sample by vortexing for 30 s and centrifugation at 3000 rpm for one minute. The supernatant was discarded, and DNA was extracted from the fecal pellet using the Norgen Stool DNA Isolation Kit, Norgen Biotek Corp (Thorold, ON, Canada), following the manufacturer's recommendations.

Blastocystis detection in animal samples followed the same methodology as previously described for human samples (Higuera et al., 2020b). The amplification of the rRNA-18S gene fragment was achieved by PCR, with a final volume of which contained 0.28X (3.5 μL) GoTaq Green Master Mix (Promega[®], Madison, Wisconsin, USA), 2 μL of template DNA, and 1 μM of each primer with a final volume of 9 μL . The primers used were Blasto FWD F5 (5'-GGTCCGGTGAACACT TTGGATTT-3') and Blasto R F2 (5'-CCTACGGAAACCTTGTACGA CTTCA-3') (Christen Rune Stensvold et al., 2012). Thermal cycling starts with 95 $^{\circ}\text{C}$ for 5 min, then 35 cycles of 95 $^{\circ}\text{C}$ for 15 s, 58 $^{\circ}\text{C}$ for 1 min, and 72 $^{\circ}\text{C}$ for 30 s; 72 $^{\circ}\text{C}$ for 10 min. The size of each amplicon was evaluated by 2% agarose gel electrophoresis stained with SYBR[™] Safe (Invitrogen[™], Carlsbad, CA, USA (Stensvold and Clark, 2016; Zhao et al., 2017). Human samples did not undergo this procedure, as *Blastocystis* identification had been previously performed using this method in the study by Higuera and colleagues (Higuera et al., 2020b).

2.3. PCR amplification and sequencing of the full-length SSU rRNA 18S gene by Oxford Nanopore MinION

Molecular characterization of *Blastocystis* was performed on PCR-positive samples from both human and animal hosts. DNA from these

samples underwent PCR to generate amplicons of the complete rRNA-18S gene, approximately 1800 bp in length, using 1 μ M of primers (Af) SSU-F1 (5-AAC CTG GTT GAT CCT GCC AGT C-3) and (Br) SSU-R1 (5-TGA TCC TTC TGC AGG TTC ACC TAC G-3) forward and reverse, respectively. This was carried out utilizing 0.3375 U (12.5 μ L) high-fidelity proofreading polymerase contained in KAPA HiFi Hot Start Ready-mix (KAPA Biosystems, Cape Town, South Africa) with a reaction volume of 25 μ L. The thermal cycling involved initial denaturation at 98 °C for 5 min, followed by 35 amplification cycles of 20 s at 98 °C, 45 s at 60 °C, and 90 s at 72 °C. A final extension of 5 min concluded the process (Maloney et al., 2020b).

The Oxford Nanopore MinION platform was utilized for characterization, following the protocol established by Maloney et al. in 2020 (Maloney et al., 2021b, 2020b). PCR amplicons were purified using 0.5 \times AMPure XP beads (Beckman Coulter Brea, CA, USA) (Higuera et al., 2021; Maloney et al., 2021b). The SQK-LSK109 and SQK-LSK110 ligation sequencing kits from Oxford Nanopore Technologies (ONT) were employed for library preparation, following the manufacturer's instructions for PCR coding amplicons (PBAC12_9112_v110_revB_10-Nov2020). The EXP-PBC001 PCR Barcoding Kit from ONT, Oxford, UK, was used in conjunction with ligation kits for barcoding each sample. Subsequent purification steps ensured the loading of 50 or 75 fmol into the flow cell R9 version. MinKNOW v20.10.06 software facilitated the sequencing runs (Maloney et al., 2020b).

2.4. Bioinformatics analysis

Basecalling was performed using Guppy v4. Reads were visualized using UGENE v46.0 software. FASTQ reads were filtered to retain lengths between 1600 and 2100 bp nucleotides. Adapters utilized for full-length sequencing PCR were trimmed, and only reads with full-length direct and reverse adapters were retained. From the generated FASTA files, subtyping was accomplished through alignment based on the best match with full-length *Blastocystis* reference nucleotide sequences, obtained from the reference database (<http://entamoeba.lsh.tm.ac.uk/blastorefseqs.htm>, accessed on 03/02/2021) along with other accepted full-length ST sequences present in GenBank, utilizing CENTRIFUGE algorithm with the following parameters: -k 1 -min-totallen 800 -qc-filter. To validate the assigned STs, the BLAST algorithm was utilized (10 std slen" -perc_identity 95 -evaluate 0.01 -max-target_seqs 1). The acquired full-length sequences were deposited in the GenBank database with the following accession numbers SAMN37309017 - SAMN37309027.

2.5. Phylogenetic reconstruction

Phylogenetic reconstruction employed full-length reference sequences which include all currently published subtypes and full-length SSU rRNA sequences obtained in this study. Alignment was performed using the MAFFT v.7 algorithm, and the tree was rooted with *Proteromonas lacertae*, a closely related Stramenopile, as an outgroup (Maloney et al., 2020b). The maximum likelihood method was applied for phylogenetic reconstruction, under nucleotide substitution model: TIM3+*F* + R6. Support for clades was determined through 1000 bootstrapping replicates, genetic distances were calculated with Kimura 2-parameter model using IQtree (Minh et al., 2020). The Interactive Tree of Life v32 tool facilitated tree editing and visualization (Letunic and Bork, 2016). Finally, phylogenetic network was also built in SplitsTree4 (Huson and Bryant, 2006) using the Neighbor-Net method.

2.6. Frequencies and analysis of the genetic diversity *Blastocystis* subtypes

Once STs were obtained and validated for the samples herein used, using ArcGIS software, a map was generated to visualize distribution of subtype frequencies in human and animal samples. This was represented through pie charts, aligned with specific geographical locations of each

department or municipality included in the study. The geographic data was extracted from a compiled database of acquired sequences. Additionally, we constructed bar plots to illustrate mixed infection variations within each sample, considering hosts and geographical regions. To differentiate the identified subtypes, a distinct color pattern was implemented. Additionally, we utilized the alignment of the 11 concatenated sequences obtained for each *Blastocystis* subtype to calculate indices of diversity, polymorphic sites, the number of haplotypes, haplotype diversity, and conducted population structure tests such as Tajima's D, Fu and Li's D and F.

3. Results

3.1. *Blastocystis* identification

A total of 277 animal feces samples were included in this study. Out of these, 87 samples (31.4%) tested positive for *Blastocystis* through PCR in both geographic regions—Bogotá and Boyacá. Specifically, within the Boyacá region, 76 samples (27.43%) from animal hosts were positive, which correspond as follows 62.8% ($n = 22$) from dogs, 36.8% ($n = 21$) from cats, 28.9% ($n = 11$) from sheep, 52.7% ($n = 19$) from pigs, and 13.6% ($n = 3$) from zoo animals (*Odocoileus virginianus*, *Dasyprocta fuliginosa*, and *Ateles hybridus*). Additionally, 28.2% ($n = 11$) of positive samples were collected from cats in Bogotá (Table 2). This suggests a wide detection of *Blastocystis* in both geographical areas. After subjecting the 87 samples to a second PCR for amplification of the full-length 18S SSU rRNA gene, positive results were obtained for 23 samples, corresponding to pigs ($n = 15$), sheep ($n = 7$) and zoo animals ($n = 1$) belonging to *Ateles hybridus*. The presence of a distinct band in the electrophoresis gel confirmed these positive results.

A total of 111 human samples were considered, selected based on the detection of *Blastocystis* via Sanger sequencing of a 600 bp region of the SSU rRNA gene, as previously conducted by Higuera et al. (Higuera et al., 2020b). These samples exhibited a high frequency of the protist in the Caribbean region, followed by the Orinoco. Out of this group, 29 samples available (Bolívar $n = 10$ and Casanare $n = 19$) were subjected to Oxford Nanopore sequencing to obtain full-length sequences.

3.2. Phylogenetic analysis

To verify the accuracy of the identified subtypes from the obtained full-length sequences of the SSU rRNA 18S gene, a phylogenetic reconstruction was conducted following the recommended guidelines (Stensvold and Clark, 2020). The eleven full-length sequences representing various subtypes (ST1, ST2, ST3, ST5, ST10, ST14, ST15, ST21, ST24, ST25, ST26) from different samples were included in this analysis. Employing the maximum likelihood method, the phylogenetic relationship among these sequences was determined. Notably, all the sequences clustered alongside the corresponding reference full-length sequences, further affirming their validity (Fig. 2).

3.3. *Blastocystis* subtypes and intra-subtype variation

Of the 198 PCR-positive samples from both human and animal hosts, 52 were sequenced for *Blastocystis* presence, 29 belonging to human samples and 23 to animal samples as follows: pigs ($n = 15$), sheep ($n = 7$) and zoo animals ($n = 1$), yielding a total of 2670,529,445 reads. After quality control by trimming and filtering, 158,732 reads remained for further analysis and phylogenetic reconstruction. Among the 52 sequenced samples, 11 distinct *Blastocystis* subtypes were identified throughout animal and human hosts, with the exception of sample 34, which had no assigned subtypes. The most prevalent subtype was ST1 detected in 43.13% ($n = 22$) in both animal and human samples. Following closely was ST3, present in 39.21% ($n = 20$) of samples from both hosts. Other subtypes were limited to either animal or human samples. Notably, ST2 was found only in human samples (7.84%, $n = 4$),




Fig. 2. Phylogenetic Relationships from the full-length SSU rRNA gene sequences obtained in the present study, *Blastocystis* representative reference sequences from each accepted ST, and *Proteromonas lacertae* as the outgroup microorganism for rooting the tree.

while subtypes ST5, ST15, ST10, ST24, ST25, and ST26 were found exclusively in animal samples, comprising 21.56% ($n = 11$), 15.68% ($n = 8$), 11.76% ($n=6$), 11.76% ($n = 6$), 11.76% ($n = 6$), and 9.80% ($n = 5$) of the samples, respectively. ST14 and ST21 were the least frequent subtypes detected in animal samples (7.84%, $n = 4$). The total of subtype frequencies exceeds 100% given detection of mixed infections with more than one *Blastocystis* subtype co-occurring within individual samples.

Genetic diversity indices were calculated by hosts (Human, Sheep and Pig), along with the entire dataset of STs (Total), showing the number of haplotypes equaled the number of sequences analyzed, indicating that each identified subtype was represented by unique sequences. The number of segregating sites in STs detected in human and sheep samples were greater than in pig samples. This finding was confirmed by analyzing the paired comparison between haplotypes (Supplementary Figure 1a) and supported by the tree topology obtained

from the phylogenetic network constructed in SplitsTree5, in which every subtype had its own branch (Supplementary Figure 1b).

The analysis of genetic diversity and population structure in three distinct groups showed that pig samples have displayed the most pronounced genetic diversity, characterized by a π value of 0.14502, indicating a high degree of genetic variability compared to the other populations. human STs exhibits a moderate level of genetic diversity, as reflected by a π value of 0.14011. In contrast, sheep samples demonstrate a lower genetic diversity with a π value of 0.09301. In terms of population genetic structure pig subtypes had the highest R2 value of 0.5 compared with the other hosts which displays a comparatively lower degree of R2 values. In general, the populations evaluated had different levels of genetic diversity and population genetic structure (Supplementary Table 3).

3.4. Blastocystis subtypes in animal hosts

A total of 23 animal samples were analyzed by Nanopore sequencing. As previously mentioned, sample 34, originating from a pig, did not yield a successful assignment using the employed methods. Hence, 22 samples were further analyzed. Across these animal samples, sequences from 10 distinct subtypes were obtained. Remarkably, ST15, which had not been previously reported in this country, was the second most common subtype, detected in 8 out of 22 samples (36.36%). ST5 was the most prevalent subtype, found in 11 out of 22 samples (50%). In terms of lower subtype frequencies, ST10, ST24, and ST25 were detected in 6 samples (27.27%). ST26 was identified in 5 out of 22 samples (22.72%), while ST1, ST14, and ST21 were each observed in 4 out of 22 samples (18.18%). Finally, ST3 was present in 3 out of 22 samples (13.63%) (Fig. 3a). These results showed that in several samples were detected more than one subtype.

Sheep samples displayed a higher STs diversity in *Blastocystis* subtypes, with six subtypes identified: ST10, ST14, ST21, ST24, ST25, and ST26. Multiple subtypes were detected within individual samples from the same sheep host, indicating mixed infections. Most sheep samples 6/7 (85.71%) displayed mixed infections, with six out of seven samples testing positive for ST10 (85.71%) and five out of seven samples (71.42%) testing positive for ST24, ST25, and ST26. Other subtypes were detected at lower frequencies: ST21 in three out of seven samples (42.85%) and ST14 in two out of seven samples (28.57%) (Fig. 4a).

In contrast, pig samples exhibited lower diversity of STs, with four identified *Blastocystis* subtypes: ST1, ST3, ST5, and ST15. Most pig samples (64.28%) showed mixed infections, with 9 out of 14 samples displaying multiple subtypes. ST15 was the most prevalent subtype in pig samples, found in 78.57% (11 out of 14) of the samples, followed by ST15 in 57.14% (8 out of 14) of the samples. Other subtypes were detected at lower frequencies: ST3 in 28.57% of the samples (4 out of 14), and ST1 in 21.42% of the samples (3 out of 14). Additionally, only one subtype, ST1, was identified in the zoo animal *Ateles hybridus* (Fig. 4a).

3.5. Variation of Blastocystis subtypes in human hosts

The examination of human samples revealed a limited diversity in *Blastocystis* subtypes, detecting three previously reported STs: ST1, ST2, and ST3. The most prevalent subtype among the samples was ST1, constituting 62.06% of cases, followed by ST3 (51.72%), and ST2 (13.79%) (Fig. 3b). These findings align with the assignments made by Higuera et al. using Sanger sequencing.

Upon analyzing the subtype diversity of human samples across different departments, Casanare exhibited the presence of two subtypes (ST1 and ST3), indicating comparatively lower STs diversity than Bolívar, where three STs were detected (ST1, ST2, and ST3). Intriguingly, the frequencies of the most abundant subtypes in these areas appeared to differ based on geographic location. In Casanare, the two identified STs were equally prevalent, representing 68.42% of the samples ($n = 13$). In contrast, in Bolívar, ST1 emerged as the dominant subtype, observed in 50% of the samples ($n = 5$), followed by ST2 in 40% of the samples ($n = 4$), and ST3 in 30% of the samples ($n = 3$) (Fig. 2). Additionally, mixed infections were observed within this group of isolates. Among the 29 analyzed human samples, eight (27.58%) displayed co-infections with two subtypes. The combination of ST1 and ST3 was exclusively detected in Casanare (36.84%), while a mixed infection of ST1 and ST2 was identified in Bolívar (10%). The distribution of *Blastocystis* subtypes between Casanare and Bolívar demonstrated certain variations in occurrences of unique and multiple STs (Fig. 4b).

4. Discussion

The present study reports the frequency and distribution of *Blastocystis* subtypes in both animal and human hosts across various locations in Colombia, including Bogotá and Boyacá for animals, and Bolívar and Casanare for humans. Interestingly, *Blastocystis* was detected in all sampled locations, indicating a widespread distribution of the micro-eukaryote in these areas. The overall detection rate of *Blastocystis* in animal feces samples was 31.4%, with positive cases found in various animal hosts, including dogs, cats, sheep, pigs, and zoo animals. These findings reveal a lower frequency compared to previous reports in Colombia, where frequencies ranged between 67%–81.4% (Higuera




Fig. 3. Geographical Distribution of *Blastocystis* Subtypes in Colombia. a. Animal Samples. b. Human samples. Marked with different colored dots, the precise locations where the sequenced samples were collected are shown on the map of Colombia. Pie charts with the percentages of subtypes by department are included. The host and number of samples used for this study are specified in the legend.




Fig. 4. *Blastocystis* Variation between STs in animal and human hosts. In this study was performed the detection of mixed and unique infection from different *Blastocystis* subtypes. Bar plots were constructed by host to reveal the percentage of each ST detected by sample. All subtypes were assigned a color to distinguish them from each other. a. *Blastocystis* STs diversity in animal hosts. b. *Blastocystis* STs diversity in humans by department.

et al., 2021; Onder et al., 2021). However, the observed prevalence is consistent with reports from other countries in Europe and Asia, ranging from 8.9% to 43.6% (AbuOdeh et al., 2019; Chen et al., 2023; Cian et al., 2017; Danišová et al., 2022; Li et al., 2019; Naguib et al., 2022; Nemati et al., 2021; Onder et al., 2021; Rauff-Adedotun et al., 2023; Rudzińska et al., 2022; Udonsom et al., 2018). Moreover, a meta-analysis highlighted that in certain countries in Southeast Asia, the prevalence can reach 100%, depending on the animal host (Rauff-Adedotun et al., 2020). Revealing a significant variability in the prevalence of the protists at both national and global levels.

This is the first study to detect higher frequency of *Blastocystis* in companion animals such as dogs and cats in Colombia (Supplementary Table 2), as is being reported domestic animals in shelter environment are in higher risk of carrying enteropathogens (Riaux et al., 2014), we hypothesized that dogs residing in rural environments, where the likelihood contact with diverse animal species, might be a contributing factor to increase the prevalence of *Blastocystis* in dogs. Previous research have primarily focused on investigating the presence of the microeukaryote and the potential associations between dogs and urban settings, including their owners infection (Mahdavi et al., 2022; Paulos et al., 2018; Potes-Morales and Crespo-Ortiz, 2023; Riaux et al., 2014; Rudzińska et al., 2022). However, there remains insufficient studies addressing the prevalence of *Blastocystis* in dogs residing in rural areas.

While the initial screening identified a high frequency of *Blastocystis*

in dog samples, there was a specific challenge attempting to amplify the complete 18S gene. Despite, the inability to perform complete gene sequencing, the preliminary data obtained from partial gene amplification was able to identify a high prevalence of the infection within the dog population. This shows that partial gene amplification can serve as an early diagnostic tool for veterinarians, researchers, and surveillance efforts. Further research and continued surveillance are needed to explore the complex epidemiology of *Blastocystis* and its potential impact on companion animals and human populations.

Although *Blastocystis* exhibits a substantial worldwide prevalence in cattle (Cabrine-Santos et al., 2021; Higuera et al., 2021; Mokhtar and Youssef, 2018; Naguib et al., 2022; Udonsom et al., 2018), our study showed no positive results in this animal samples. This discrepancy may be attributed to various factors. Firstly, the methodology employed, including diagnosis criteria, sample selection and the chosen location from a specific geographic region of Colombia and specific point on time, may have played a role in this results. Moreover, cattle conditions can vary and contribute to variable prevalences, the unique characteristics of the cattle population and regional differences, might influence in the absence of positive cattle samples. Furthermore, gut environment and microbiota composition of cattle are complex and dynamic, due to factors such as diet, and breed. To the best of our knowledge, all of the animals included in this study have healthy conditions. However, the overall health and immune status and management practices can impact

in the susceptibility of enteric microorganism (Xu et al., 2021). The partial differences between the local experimental context and global perspective should be approach with a more extensive dataset. This could help understanding the complex interplay of factors that influenced the outcomes of our study and contribute to a more comprehensive understanding of *Blastocystis* prevalence in cattle.

Significantly, the current analysis revealed differences in subtype distribution among hosts and regions. *Blastocystis* has been identified in human samples throughout the Americas, Europe, Africa and Asia, these molecular studies have reported the presence of commonly isolated STs in humans including ST1-ST4 (Ahmed et al., 2022; Ajampur and Tan, 2016; Gabrielli et al., 2020; Jiménez et al., 2022; P.A. 2019; Munsaka et al., 2022; Poulsen et al., 2016; Stensvold et al., 2020). In Colombia, subtypes ST1 to ST3 were identified previously. Particularly, ST2 in the Bolivar region (Higuera et al., 2020b), which was exclusively detected in human samples from Bolivar in our study, suggesting a potential host-specific preference for this subtype and regional variation in *Blastocystis* prevalence and subtype frequency (Fig. 3b) (Stensvold and Clark, 2020). In contrast, several subtypes (ST5, ST10, ST14, ST15, ST21, ST24, ST25 and ST26) were mainly found in animal samples. The analysis of samples from animals revealed a higher variety of *Blastocystis* subtypes in comparison to the human samples (Supplementary Table 3). This subtype variety may occurred due to various factors, such as interactions between different animal species, their environments, and their immune responses to the microorganism (Alzate et al., 2020; Deng et al., 2023, 2021; Iebba et al., 2016; Rojas-Velázquez et al., 2022; Yañez et al., 2021). Notably, our results revealed an absence of overlapping *Blastocystis* subtypes between different animal species, such as pigs and sheep (Supplementary Table 1) (Supplementary Figure 1). Instead, distinct subtypes were observed in each group, suggesting unique subtype assignments for each animal category (Fig. 3a). This pattern may suggest that certain *Blastocystis* subtypes have a specific host tendency, potentially pointing towards divergent transmission routes for these subtypes among animals within the same geographical area (Deng et al., 2021; Jinatham et al., 2021).

In addition to the above results, *Blastocystis* subtyping detected a total of four different subtypes in pig samples, among which ST5 and ST15 were the most common (Fig. 4a). The high prevalence of *Blastocystis* in pigs has been previously associated with host factors such as immune system status, gender, and age, the latter being the principal property which had been related to its prevalence (Hublin et al., 2020). Remarkably, ST15 was detected for the first time in Colombia during this study. It has been suggested that ST15 has a mammalian origin, the elevated detection of ST15 in pig feces supports this hypothesis (Daníšová and Valenčáková, 2021; Wylezich et al., 2019), and previously reported in camels, buffalo, wild boars and gibbons (Asghari et al., 2021; Pawestri et al., 2022; Russini et al., 2020), as it has been identified most in domesticated animals and livestock, which are in close proximity to humans; however, its presence has not been confirmed in humans, further research should be needed to determine its potential for zoonotic transmission (Supplementary Table 4). The identification of subtypes, such as ST15 in animals, highlights the use of Next-Generation Sequencing (NGS) technology to explore the diversity of *Blastocystis* subtypes, the importance of continuous surveillance to monitor the emergence of undetected subtypes and their potential implications for public health.

Blastocystis had been detected in a wide host range including mammals, birds, reptiles and mollusks worldwide (Supplementary Table 4), this broad host range suggest that animals might serve as reservoirs of the microeukaryote, the contact with infected animals, their feces or contaminated environment had been linked to human transmission (Hernández et al., 2023; Rudzińska and Sikorska, 2023; Stensvold and Clark, 2016). For instance, livestock, pets and wildlife animals which has close contact with humans such as keepers, owners, veterinarians or farmers, along with contact with other animal species may play an important role in the transmission and serve as potential reservoir for

other domestic animals and humans infections (Naguib et al., 2022; Salehi et al., 2022). In this study companion pets such as cats and dogs showed a moderate frequency of *Blastocystis*, this proximity could increase the risk of zoonotic transmission, particularly if pets carry possible zoonotic subtypes. Recently, Hernández et al. demonstrated variable associations between *Blastocystis* infection in Colombian schoolchildren and contact with domestic animals. Comparisons between infection status and animal exposure revealed increased odds of ST1 and mixed subtype infections with dogs and cats living in the same house. Conversely, the presence of chickens was associated with decreased odds for multiple *Blastocystis* infection outcomes. The dichotomous risks observed with companion animals compared to chickens highlight that animal reservoirs and zoonotic transmission dynamics of *Blastocystis* can be complex (Hernández et al., 2023). Additionally, considering companion animal owners can harbor *Blastocystis* without showing clinical signs, potentially exposing other humans or animals to the protist. Further investigations utilizing molecular epidemiology approaches are warranted to clarify potential zoonotic and animal-human transmission routes for this common intestinal parasite. Elucidating the relationships between *Blastocystis* genetic diversity across host species will provide crucial insights into its public health significance. The detection of *Blastocystis* in diverse animal species indicates the potential for zoonotic transmission and the need for further investigation to understand the transmission dynamics between animals and humans (Maloney et al., 2021a).

Among the animal hosts, sheep exhibited the highest diversity of *Blastocystis* subtypes, with seven different subtypes identified which correspond to previously reported subtypes in the same region (Higuera et al., 2021) (Supplementary Table 4). A considerable proportion of sheep samples showed evidence of mixed infections, suggesting a tendency for co-infection with multiple *Blastocystis* subtypes within individual hosts. Moreover, mixed infections were observed in some human samples, with co-occurrence of two subtypes (ST1 and ST3) although less frequently compared to animals (Fig. 4). The presence of multiple subtypes in a single host may be influenced by various factors, including alterations in the gut microbiota, which can involve the presence of diverse bacterial species and changes in the microenvironment (Vega et al., 2021; Vega Romero, 2020). Host immune responses, as well as factors related to food and water supplies, also likely contribute to shaping the diversity of microorganisms, including *Blastocystis* (Deng et al., 2023; Jinatham et al., 2021). Socioeconomic factors, such as poverty or inadequate living conditions, may further contribute to the acquisition of gut parasitism, as individuals may be more exposed to such infections (Higuera et al., 2020b). These findings suggest possible interactions and competition between *Blastocystis* subtypes (Supplementary Figure 1). However, further studies are required to determine whether these interactions follow specific patterns that facilitate the coexistence of multiple subtypes within a single host. Knowledge of these mechanisms may lead to a better understanding on the maintenance and prevalence of *Blastocystis* subtypes in animal populations.

In the present study we identified the presence of *Blastocystis* subtypes ST1 and ST3 in both human and animal hosts. These subtypes have previously been linked to gastrointestinal symptoms such as diarrhea, abdominal pain, and urticaria in humans, and there have been suggestions of an association with Inflammatory Bowel Disease (IBD) (Peña et al., 2020; S. 2019). Interestingly, we observed the presence of these subtypes in pigs and a monkey from the zoo, which are in regular contact with their keepers, farmers in the case of pigs, and zookeepers in the case of the monkey (Supplementary Table 2). This close human-animal interaction poses a potential risk of zoonotic transmission, whereby *Blastocystis* may be transmitted between animals and humans (Asghari et al., 2021; Cian et al., 2017; Naguib et al., 2022; Rauff-Adedotun et al., 2023; Stensvold and Clark, 2020). Such transmission may result in the dissemination of pathogenic subtypes and could have implications for public health and the welfare of both animals and humans.

While this study provides valuable information on *Blastocystis*

frequency and subtype distribution in some regions from Colombia, it also has several limitations that should be considered. Firstly, the used samples may not represent the entire population of humans and animal species in the sampled regions, even though is informative, a larger sample size would enhance statistical robustness. Moreover, due to the cross-sectional design, this study shows the prevalence of *Blastocystis* on a specific point in time which makes it impossible to track the changes of the microeukaryote over the time, demanding further longitudinal investigations. Furthermore, this study lacks detailed clinical data from human hosts restricting the understanding of potential associations between the subtypes and clinical symptoms. Although this study suggests the potential for zoonotic transmission of *Blastocystis*, it does not directly investigate transmission dynamics between animals and humans or public health risks; additional investigation including contact tracing and extend human health impact is critical for confirmation of transmission and exploring factors influencing health implications. Furthermore, in this study it is important to consider the relatively high error rate introduced in sequences using nanopore technology, which can be attributed to base-calling inaccuracies, overestimated quality scores, errors in homopolymeric or short repeat regions (Delahaye and Nicolas, 2021; Sahlin and Medvedev, 2021), it is crucial to recognize that despite of ensuring updated basecaller and executing manual base-calling algorithm to improve accuracy and mitigate the impact in the obtained results, this technology continues to be improved and may introduce errors.

5. Conclusion

Finally, the present study demonstrates that *Blastocystis* is prevalent in both humans and animals, and the use of NGS allows for the identification of a variety of subtypes, showing both inter and intra-subtype variation. The implementation of Next-Generation Sequencing (NGS) proved instrumental in providing a thorough molecular characterization of *Blastocystis*. The identification of suspected zoonotic subtypes provides valuable information to understand the transmission dynamics of the parasite and potential role of animals as reservoirs for *Blastocystis* subtypes. However, future studies should consider other regions of the country, different domestic and wild animal species, and variables describing the health status of the individuals to determine potential influences on parasite presence. Additionally, establishing relationships between humans and animals in contact with each other is crucial for a comprehensive understanding of *Blastocystis* transmission pathways. A One Health approach, involving collaboration between human and animal health sectors, is essential to address the potential public health implications of *Blastocystis* transmission and establish effective preventive strategies.

Author contributions

LCS, AC, collected the samples. AC,LS,PAJ made the DNA extraction. PAJ performed the PCRs. PAJ and LCS performed NGS sequencing with MinION. PAJ analyzed the data and wrote the manuscript. PAJ and MM reviewed and edited the manuscript. JDR conceived the manuscript, supervised the project, wrote, reviewed, and edited the manuscript. All authors approved the final version of the manuscript.

CRedit authorship contribution statement

Paula Jiménez: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Resources, Supervision, Validation, Visualization, Writing – original draft. **Marina Muñoz:** Conceptualization, Data curation, Formal analysis, Resources, Supervision, Writing – review & editing. **Lissa Cruz-Saavedra:** Data curation, Formal analysis, Resources, Writing – review & editing. **Anny Camargo:** Investigation, Methodology, Resources, Writing – review & editing. **Juan David Ramírez:** Conceptualization, Data curation, Formal analysis, Funding

acquisition, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing – original draft.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgments

This study was funded by Dirección de Investigación e Innovación from Universidad del Rosario.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.actatropica.2023.107090.

References

- AbuOdeh, R., Ezzedine, S., Madkour, M., Stensvold, C.R., Samie, A., Nasrallah, G., AlAbsi, E., ElBakri, A., 2019. Molecular Subtyping of *Blastocystis* from Diverse Animals in the United Arab Emirates. *Protist* 170, 125679. <https://doi.org/10.1016/j.protis.2019.125679>.
- Ahmed, S.A., El-Mahallawy, H.S., Mohamed, S.F., Angelici, M.C., Hasapis, K., Saber, T., Karanis, P., 2022. Subtypes and phylogenetic analysis of *Blastocystis* sp. isolates from West Ismailia. *Sci. Rep.* 12, 19084. <https://doi.org/10.1038/s41598-022-23360-0>.
- Ajjampur, S.S., Tan, K.S., 2016. Pathogenic mechanisms in *Blastocystis* spp.— Interpreting results from in vitro and in vivo studies. *Parasitol. Int.*
- Alfellani, M.A., Stensvold, C.R., Vidal-Lapedra, A., Onuoha, E.S.U., Fagbenro-Beyioku, A.F., Clark, C.G., 2013a. Variable geographic distribution of *Blastocystis* subtypes and its potential implications. *Acta Trop.*
- Alfellani, M.A., Taner-Mulla, D., Jacob, A.S., Imeede, C.A., Yoshikawa, H., Stensvold, C.R., Clark, C.G., 2013b. Genetic diversity of *Blastocystis* in livestock and zoo animals. *Protist*.
- Alzate, J.F., Toro-Londoño, M., Cabarcas, F., Garcia-Montoya, G., Galvan-Diaz, A., 2020. Contrasting microbiota profiles observed in children carrying either *Blastocystis* spp. or the commensal amoebas *Entamoeba coli* or *Endolimax nana*. *Sci. Rep.* 10, 15354. <https://doi.org/10.1038/s41598-020-72286-y>.
- Asghari, A., Sadrebazazz, A., Shamsi, L., Shams, M., 2021. Global prevalence, subtypes distribution, zoonotic potential, and associated risk factors of *Blastocystis* sp. in domestic pigs (*Sus domesticus*) and wild boars (*Sus scrofa*): A systematic review and meta-analysis. *Microb. Pathog.* 160, 105183 <https://doi.org/10.1016/j.micpath.2021.105183>.
- Baek, S., Maloney, J.G., Molokin, A., George, N.S., Cortés Vecino, J.A., Santin, M., 2022. Diversity of *Blastocystis* Subtypes in Horses in Colombia and Identification of Two New Subtypes. *Microorganisms* 10, 1693. <https://doi.org/10.3390/microorganisms10091693>.
- Barbosa, C.V., Batista, R., de J., Igreja, R.P., Levy, C.M.d'Avila, Werneck de Macedo, H., Santos, H.L.C., 2017. Distribution of *Blastocystis* subtypes isolated from humans from an urban community in Rio de Janeiro, Brazil. *Parasites & Vectors* 10 (1), 1–9. <https://doi.org/10.1186/s13071-017-2458-0>. Iss2017.
- Bowden, R., Davies, R.W., Heger, A., Pagnamenta, A.T., de Cesare, M., Oikkonen, L.E., Parkes, D., Freeman, C., Dhalla, F., Patel, S.Y., Popitsch, N., Ip, C.L.C., Roberts, H.E., Salatino, S., Lockstone, H., Lunter, G., Taylor, J.C., Buck, D., Simpson, M.A., Donnelly, P., 2019. Sequencing of human genomes with nanopore technology. *Nat. Commun.* 10, 1869. <https://doi.org/10.1038/s41467-019-09637-5>.
- Cabrine-Santos, M., Moura, R.G.F., Pedrosa, A.L., Correia, D., Oliveira-Silva, M.B.de, 2021. Molecular characterization of *Blastocystis* subtypes isolated in the city of Uberaba, Minas Gerais State, Brazil. *Rev. Soc. Bras. Med. Trop.* 54 <https://doi.org/10.1590/0037-8682-0305-2021>.
- Casero, R.D., Mongi, F., Sánchez, A., Ramírez, J.D., 2015. *Blastocystis* and urticaria: Examination of subtypes and morphotypes in an unusual clinical manifestation. *Acta Trop.* <https://doi.org/10.1016/j.actatropica.2015.05.004>.
- Castañeda, S., Muñoz, M., Villamizar, X., Hernández, P.C., Vásquez, L.R., Tito, R.Y., Ramírez, J.D., 2020. Microbiota characterization in *Blastocystis*-colonized and *Blastocystis*-free school-age children from Colombia. *Parasites & Vectors* 13, 521. <https://doi.org/10.1186/s13071-020-04392-9>.
- Chen, H., Hao, Y., Liu, Y., Xu, M., Zhang, W., Li, H., Yang, F., 2023. The frequency and subtype distribution of *Blastocystis* sp. in humans and domestic animals in

- households in Heilongjiang Province, China. *Acta Trop.* 240, 106844 <https://doi.org/10.1016/j.actatropica.2023.106844>.
- Cian, A., El Safadi, D., Osman, M., Moriniere, R., Gantois, N., Benamrouz-Vanneste, S., Delgado-Viscogliosi, P., Guyot, K., Li, L.L., Monchy, S., 2017. Molecular epidemiology of *Blastocystis* sp. in various animal groups from two French zoos and evaluation of potential zoonotic risk. *PLoS One*.
- Clark, C.G., Van der Giezen, M., Alfellani, M.A., Stensvold, C.R., 2013. Recent developments in *Blastocystis* research.
- Cruz-Saavedra, L., Arévalo, V.A., García-Corredor, D., Jiménez, P.A., Vega, L., Pulido-Medellín, M., Ortiz-Pineda, M., Ramírez, J.D., 2023. Molecular detection and characterization of *Giardia* spp., *Cryptosporidium* spp., and *Blastocystis* in captive wild animals rescued from central Colombia. *Int. J. Parasitol. Parasites Wildl.* 22, 1–5. <https://doi.org/10.1016/j.ijppaw.2023.07.005>.
- Danišová, O., Valenčíková, A., 2021. First detection of *Blastocystis* sp. in pigs in Slovakia and in Europe. *Parasitol. Int.* 81, 102235 <https://doi.org/10.1016/j.parint.2020.102235>.
- Danišová, O., Valenčíková, A., Kandrácová, P., Tomko, M., Sučík, M., 2022. First report of *Blastocystis* spp. subtypes in ZOO animals in Slovakia, Central Europe. *Ann. Agric. Environ. Med.* 29, 149–151. <https://doi.org/10.2644/aaem/145826>.
- Delahaye, C., Nicolas, J., 2021. Sequencing DNA with nanopores: Troubles and biases. *PLoS One* 16, e0257521. <https://doi.org/10.1371/journal.pone.0257521>.
- Deng, L., Wojciech, L., Gascoigne, N.R., Peng, G., Tan, K.S., 2021. New insights into the interactions between *Blastocystis*, the gut microbiota, and host immunity. *PLoS Pathog.*
- Deng, L., Wojciech, L., Png, C.W., Kioh, D.Y.Q., Gu, Y., Aung, T.T., Malleret, B., Chan, E. C.Y., Peng, G., Zhang, Y., Gascoigne, N.R.J., Tan, K.S.W., 2023. Colonization with two different *Blastocystis* subtypes in DSS-induced colitis mice is associated with strikingly different microbiome and pathogenic features. *Theranostics* 13, 1165–1179. <https://doi.org/10.7150/tno.81583>.
- Gabrielli, S., Furzi, F., Fontanelli Sulekova, L., Taliani, G., Mattiucci, S., 2020. Occurrence of *Blastocystis*-subtypes in patients from Italy revealed association of ST3 with a healthy gut microbiota. *Parasite Epidemiol. Control.* <https://doi.org/10.1016/j.parepi.2020.e00134>.
- Hernández, P.C., Maloney, J.G., Molokin, A., George, N.S., Morales, L., Chaparro-Olaya, J., Santin, M., 2023. Exploring *Blastocystis* genetic diversity in rural schoolchildren from Colombia using next-generation amplicon sequencing reveals significant associations between contact with animals and infection risk. *Parasitol. Res.* 122, 1451–1462. <https://doi.org/10.1007/s00436-023-07841-3>.
- Hernández-Castro, C., Maloney, J.G., Agudelo-López, S.P., Toro-Londoño, M.A., Botero-Garcés, J.H., Orozco, M.C., Quintero-Quinchía, Y.C., Correa-Cote, J.C., Múnera-Duque, A., Ricaurte-Ciro, J.C., Londoño-Alvarez, L.L., Escobar, R.M., Köster, P.C., Sánchez, S., Carmenta, D., Santín, M., 2023. Identification and validation of novel *Blastocystis* subtype ST41 in a Colombian patient undergoing colorectal cancer screening. *J. Eukaryot. Microbiol.* 70, e12978. <https://doi.org/10.1111/jeu.12978>.
- Higuera, A., Herrera, G., Jimenez, P., García-Corredor, D., Pulido-Medellín, M., Bulla-Castañeda, D.M., Pinilla, J.C., Moreno-Pérez, D.A., Maloney, J.G., Santín, M., 2021. Identification of multiple *Blastocystis* subtypes in domestic animals from Colombia using amplicon-based next generation sequencing. *Front. Vet. Sci.*
- Higuera, A., Muñoz, M., López, M.C., Reyes, P., Urbano, P., Villalobos, O., Ramírez, J.D., 2020a. Succinate dehydrogenase gene as a marker for studying *Blastocystis* genetic diversity. *Heliyon*.
- Higuera, A., Villamizar, X., Herrera, G., Giraldo, J.C., Vasquez-A, L.R., Urbano, P., Villalobos, O., Tovar, C., Ramírez, J.D., 2020b. Molecular detection and genotyping of intestinal protozoa from different biogeographical regions of Colombia. *PeerJ* 8, e8554. <https://doi.org/10.7717/peerj.8554>.
- Hublin, J.S., Maloney, J.G., Santin, M., 2020. *Blastocystis* in domesticated and wild mammals and birds. *Res. Vet. Sci.*
- Huson, D.H., Bryant, D., 2006. Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* 23, 254–267. <https://doi.org/10.1093/molbev/msj030>.
- Iebba, V., Santangelo, F., Totino, V., Pantanella, F., Monsia, A., Di Cristanziano, V., Di Cave, D., Schippa, S., Berrilli, F., D'Alfonso, R., 2016. Gut microbiota related to *Giardia* duodenalis, *Entamoeba* spp. and *Blastocystis hominis* infections in humans from Côte d'Ivoire. *J. Infect. Dev. Ctries.* <https://doi.org/10.3855/jidc.8179>.
- Jain, M., Koren, S., Miga, K.H., Quick, J., Rand, A.C., Sasani, T.A., Tyson, J.R., Beggs, A. D., Dilthey, A.T., Fiddes, I.T., Malla, S., Marriott, H., Nieto, T., O'Grady, J., Olsen, H. E., Pedersen, B.S., Rhie, A., Richardson, H., Quinlan, A.R., Snutch, T.P., Tee, L., Paten, B., Phillippy, A.M., Simpson, J.T., Loman, N.J., Loose, M., 2018. Nanopore sequencing and assembly of a human genome with ultra-long reads. *Nat. Biotechnol.* 36, 338–345. <https://doi.org/10.1038/nbt.4060>.
- Jiménez, P., Muñoz, M., Ramírez, J.D., 2022. An update on the distribution of *Blastocystis* subtypes in the Americas. *Heliyon*, e12592.
- Jiménez, P.A., Jaimes, J.E., Ramírez, J.D., 2019. A summary of *Blastocystis* subtypes in North and South America. *Parasites & vectors*.
- Jinatham, V., Maxamhud, S., Popluechai, S., Tsaousis, A.D., Gentekaki, E., 2021. *Blastocystis* One Health Approach in a Rural Community of Northern Thailand: Prevalence, Subtypes and Novel Transmission Routes. *Front. Microbiol.* 12.
- Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* 30, 772–780. <https://doi.org/10.1093/molbev/mst010>.
- Lee, L., Chye, T., Karmacharya, B., Govind, S., 2012. *Blastocystis* sp.: waterborne zoonotic organism, a possibility? *Parasites & Vectors* 5 (1), 130. <https://doi.org/10.1186/1756-3305-5-130>. Iss2012.
- Letunic, I., Bork, P., 2016. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic. Acids. Res.* 44, W242–W245. <https://doi.org/10.1093/nar/gkw290>.
- Li, J., Karim, Md.R., Li, D., Rahaman Sumon, S.M.M., Siddiki, S.H.M.F., Rume, F.I., Sun, R., Jia, Y., Zhang, L., 2019. Molecular characterization of *Blastocystis* sp. in captive wildlife in Bangladesh National Zoo: Non-human primates with high prevalence and zoonotic significance. *Int. J. Parasitol. Parasites Wildl.* 10, 314–320. <https://doi.org/10.1016/j.ijppaw.2019.11.003>.
- Li, L.H., Zhou, X.N., Du, Z.W., Wang, X.Z., Wang, L.B., Jiang, J.Y., Yoshikawa, H., Steinmann, P., Utzinger, J., Wu, Z., Chen, J.X., Chen, S.H., Zhang, L., 2007. Molecular epidemiology of human *Blastocystis* in a village in Yunnan province, China. *Parasitol. Int.*
- Londoño-Franco, Á.L., Loaiza-Herrera, J., Lora-Suárez, F.M., Gómez-Marín, J.E., 2014. Frecuencia y fuentes de *Blastocystis* sp. en niños de 0 a 5 años de edad atendidos en hogares infantiles públicos de la zona urbana de Calarcá, Colombia /*Blastocystis* sp. frequency and sources among children from 0 to 5 years of age attending public day care centers in Calarcá, Colombia. *Biomedica.* <https://doi.org/10.7705/biomedica.v34i2.1214>.
- Mahdavi, F., Asghari, A., Shahabi, S., Shamsi, L., Soltani-Jazi, F., Sadrebazzaz, A., Shams, M., 2022. Distribution, genetic diversity, and zoonotic significance of *Blastocystis* subtypes in pet dogs. *Comp. Immunol. Microbiol. Infect. Dis.* 88, 101848 <https://doi.org/10.1016/j.cimid.2022.101848>.
- Maloney, J.G., da Cunha, M.J., Molokin, A., Cury, M.C., Santin, M., 2021a. Next-generation sequencing reveals wide genetic diversity of *Blastocystis* subtypes in chickens including potentially zoonotic subtypes. *Parasitol. Res.*
- Maloney, J.G., Jang, Y., Molokin, A., George, N.S., Santin, M., 2021b. Wide Genetic Diversity of *Blastocystis* in White-Tailed Deer (*Odocoileus virginianus*) from Maryland, USA. *Microorganisms*.
- Maloney, J.G., Lombard, J.E., Urie, N.J., Shively, C.B., Santin, M., 2018. Zoonotic and genetically diverse subtypes of *Blastocystis* in US pre-weaned dairy heifer calves. *Parasitol. Res.* <https://doi.org/10.1007/s00436-018-6149-3>.
- Maloney, J.G., Molokin, A., da Cunha, M.J.R., Cury, M.C., Santin, M., 2020a. *Blastocystis* subtype distribution in domestic and captive wild bird species from Brazil using next generation amplicon sequencing. *Parasite Epidemiol. Control.*
- Maloney, J.G., Molokin, A., Santin, M., 2020b. Use of Oxford Nanopore MinION to generate full-length sequences of the *Blastocystis* small subunit (SSU) rRNA gene. *Parasites & Vectors*.
- Maloney, J.G., Molokin, A., Santin, M., 2019. Next generation amplicon sequencing improves detection of *Blastocystis* mixed subtype infections. *Infection. Genetics and Evolution*.
- Maloney, J.G., Molokin, A., Seguí, R., Maravilla, P., Martínez-Hernández, F., Villalobos, G., Tsaousis, A.D., Gentekaki, E., Muñoz-Antolí, C., Klisiowicz, D.R., Oishi, C.Y., Toledo, R., Esteban, J.G., Köster, P.C., de Lucio, A., Dashti, A., Bailo, B., Calero-Bernal, R., González-Barrio, D., Carmenta, D., Santín, M., 2023. Identification and Molecular Characterization of Four New *Blastocystis* Subtypes Designated ST35-ST38. *Microorganisms* 11, 46. <https://doi.org/10.3390/microorganisms11010046>.
- Maloney, J.G., Santin, M., 2021. Mind the gap: New full-length sequences of *Blastocystis* subtypes generated via Oxford Nanopore Minion sequencing allow for comparisons between full-length and partial sequences of the small subunit of the ribosomal RNA gene. *Microorganisms*.
- Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear, R., 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* 37, 1530–1534. <https://doi.org/10.1093/molbev/msaa015>.
- Ministerio de Salud y Protección Social, 2015. Encuesta Nacional de Parasitismo Intestinal en Población Escolar, Colombia 2012 - 2014.
- Mokhtar, A., Youssef, A., 2018. Subtype analysis of *Blastocystis* spp. isolated from domestic mammals and poultry and its relation to transmission to their in-contact humans in Ismailia governorate, Egypt. *Parasitol. United J.* 11, 90–98.
- Munsaka, G., Hayashida, K., Mubemba, B., Simulundu, E., Mulunda, N., Pule, R., Sianongo, S., Makuluni, M., Muleya, W., Changula, K., Chitanga, S., Mutengo, M., 2022. Molecular subtyping of *Blastocystis* sp. detected in patients at a large tertiary referral hospital in Lusaka, Zambia. *Front. Parasitol.* 1.
- Naguib, D., Gantois, N., Desramaut, J., Arafat, N., Even, G., Certad, G., Chabé, M., Viscogliosi, E., 2022. Prevalence, Subtype Distribution and Zoonotic Significance of *Blastocystis* sp. Isolates from Poultry, Cattle and Pets in Northern Egypt. *Microorganisms* 10, 2259. <https://doi.org/10.3390/microorganisms10112259>.
- Nemati, S., Zali, M.R., Johnson, P., Mirjalali, H., Karanis, P., 2021. Molecular prevalence and subtype distribution of *Blastocystis* sp. in Asia and in Australia. *J. Water Health* 19, 687–704. <https://doi.org/10.2166/wh.2021.011>.
- Onder, Z., Yildirim, A., Pekmezci, D., Duzlu, O., Pekmezci, G.Z., Ciloglu, A., Simsek, E., Kokcu, N.D., Yetismis, G., Ercan, N., Inci, A., 2021. Molecular identification and subtype distribution of *Blastocystis* sp. in farm and pet animals in Turkey. *Acta Trop.* 220, 105939 <https://doi.org/10.1016/j.actatropica.2021.105939>.
- Osorio-Pulgarin, M.I., Higuera, A., Beltran-Alzate, J.C., Sánchez-Jiménez, M., Ramírez, J. D., 2021. Epidemiological and Molecular Characterization of *Blastocystis* Infection in Children Attending Daycare Centers in Medellín, Colombia. *Biology* 10, 669. <https://doi.org/10.3390/biology10070669>.
- Paulos, S., Köster, P.C., de Lucio, A., Hernández-de-Mingo, M., Cardona, G.A., Fernández-Crespo, J.C., Stensvold, C.R., Carmenta, D., 2018. Occurrence and subtype distribution of *Blastocystis* sp. in humans, dogs and cats sharing household in northern Spain and assessment of zoonotic transmission risk. *Zoonoses Public Health* 65, 993–1002. <https://doi.org/10.1111/zph.12522>.
- Pawestri, A.R., Padungsombat, J., Deesithivech, O., Nakayama, E., Shioda, T., Moonsom, S., 2022. *Blastocystis* spp. Subtype Distribution from Human and Animals at the Thai-Myanmar Border: The Public Health Implication. *Int. International Journal of Infectious Diseases, Abstracts from the Eighth International Meeting on Emerging Diseases and Surveillance, IMED 2021*, 116, p. S80. <https://doi.org/10.1016/j.ijid.2021.12.190>.

- Peña, S., Carrasco, G., Rojas, P., Castillo, D., Ozaki, L.S., Mercado, R., 2020. Determination of subtypes of *Blastocystis* sp. in Chilean patients with and without inflammatory bowel syndrome. A preliminary report. *Parasite Epidemiol. Control*. <https://doi.org/10.1016/j.parepi.2019.e00125>.
- Peña, S., Carrasco, G., Rojas, P., Gotteland, M., Castillo, D., Ozaki, L.S., Mercado, R., 2019. Different Subtypes of *Blastocystis hominis* (ST1, ST2 and ST4) are present in Chilean patients with Inflammatory Bowel Disease.
- Potes-Morales, C., Crespo-Ortiz, M.del P., 2023. Molecular diagnosis of intestinal protozoa in young adults and their pets in Colombia. *PLoS One* 18, e0283824. <https://doi.org/10.1371/journal.pone.0283824>.
- Poulsen, C.S., Efunshile, A.M., Nelson, J.A., Stensvold, C.R., 2016. Epidemiological Aspects of *Blastocystis* Colonization in Children in Ilero, Nigeria. *Am. J. Trop. Med. Hyg.* 95, 175–179. <https://doi.org/10.4269/ajtmh.16-0074>.
- Qi, M., Wei, Z., Zhang, Y., Zhang, Q., Li, J., Zhang, L., Wang, R., 2020. Genetic diversity of *Blastocystis* in kindergarten children in southern Xinjiang, China. *Parasites & Vectors*.
- Ramírez, J.D., Flórez, C., Olivera, M., Bernal, M.C., Giraldo, J.C., 2017. *Blastocystis* subtyping and its association with intestinal parasites in children from different geographical regions of Colombia. *PLoS One*. <https://doi.org/10.1371/journal.pone.0172586>.
- Ramirez, J.D., Sanchez, A., Hernandez, C., Florez, C., Consuelo Bernal, M., Cesar Giraldo, J., Reyes, P., Consuelo Lopez, M., Garcia, L., Cooper, P.J., Vicuna, Y., Mongi, F., Casero, R.D., 2016. Geographic distribution of human *Blastocystis* subtypes in South America. [10.1016/j.meegid.2016.03.017](https://doi.org/10.1016/j.meegid.2016.03.017).
- Rauff-Adedotun, A.A., Lee, I.L., Abd Talib, N., Shaari, N., Yahaya, Z.S., Meor Termizi, F. H., 2023. Prevalence, potential risk factors and genetic diversity of *Blastocystis* in ruminant livestock animals from Penang, Malaysia. *Parasitol. Res.* <https://doi.org/10.1007/s00436-023-07920-5>.
- Rauff-Adedotun, A.A., Mohd Zain, S.N., Farah Haziqah, M.T., 2020. Current status of *Blastocystis* sp. in animals from Southeast Asia: a review. *Parasitol. Res.* 119, 3559–3570. <https://doi.org/10.1007/s00436-020-06828-8>.
- Roberts, T., Stark, D., Harkness, J., Ellis, J., 2013. Subtype distribution of *Blastocystis* isolates from a variety of animals from New South Wales, Australia. *Vet. Parasitol.*
- Rojas-Velázquez, L., Morán, P., Serrano-Vázquez, A., Portillo-Bobadilla, T., González, E., Pérez-Juárez, H., Hernández, E., Partida-Rodríguez, O., Nieves-Ramírez, M., Padilla, A., Zaragoza, M., Kiménez, C., 2022. The regulatory function of *Blastocystis* spp. on the immune inflammatory response in the gut microbiome. *Front. Cell. Infect. Microbiol.* 12.
- Rondón, S., Cavallero, S., Link, A., González, C., D'Amelio, S., 2023. Prevalence and Molecular Characterisation of *Blastocystis* sp. Infecting Free-Ranging Primates in Colombia. *Pathogens* 12, 569. <https://doi.org/10.3390/pathogens12040569>.
- Ruau, G., Craig, Stang, B., 2014. Prevalence of *Blastocystis* in shelter-resident and client-owned companion animals in the US Pacific Northwest. *PLoS One* 9 (9), e107496. <https://doi.org/10.1371/journal.pone.0107496>. Iss2014.
- Rudzińska, M., Kowalewska, B., Kurpas, M., Szostakowska, B., 2022. Rare Occurrence of *Blastocystis* in Pet Animals and Their Owners in the Pomeranian Voivodeship in Poland in the Light of Literature Data. *J. Clin. Med.* 11, 2975. <https://doi.org/10.3390/jcm11112975>.
- Rudzińska, M., Sikorska, K., 2023. Epidemiology of *Blastocystis* Infection: A Review of Data from Poland in Relation to Other Reports. *Pathogens* 12, 1050. <https://doi.org/10.3390/pathogens12081050>.
- Russini, V., Di Filippo, M.M., Fanelli, R., Polidori, M., Berrilli, F., Di Cave, D., Novelletto, A., Calderini, P., 2020. Characterization of prevalence and genetic subtypes of *Blastocystis* sp. in wild and domestic Suidae of central Italy aided by amplicon NGS. *Veterinary Parasitology: Regional Studies and Reports* 22, 100472. <https://doi.org/10.1016/j.vprsr.2020.100472>.
- Sahlin, K., Medvedev, P., 2021. Error correction enables use of Oxford Nanopore technology for reference-free transcriptome analysis. *Nat. Commun.* 12, 2. <https://doi.org/10.1038/s41467-020-20340-8>.
- Salehi, R., Rostami, A., Mirjalali, H., Stensvold, C.R., Haghighi, A., 2022. Genetic characterization of *Blastocystis* from poultry, livestock animals and humans in the southwest region of Iran—Zoonotic implications. *Transbound. Emerg. Dis.* 69, 1178–1185.
- Santin, M., Figueiredo, A., Molokin, A., George, N.S., Köster, P.C., Dashti, A., González-Barrio, D., Carmena, D., Maloney, J.G., n.d. Division of *Blastocystis* ST10 into three new subtypes: ST42-ST44. *J. Eukaryot. Microbiol.* n/a, e12998. [10.1111/jeu.12998](https://doi.org/10.1111/jeu.12998).
- Stensvold, C.R., 2013. *Blastocystis*: Genetic diversity and molecular methods for diagnosis and epidemiology. *Trop. Parasitol.* <https://doi.org/10.4103/2229-5070.113896>.
- Stensvold, Christen Rune, Ahmed, U.N., Andersen, L.O., Nielsen, H.V., 2012. Development and evaluation of a genus-specific, probe-based, internal-process-controlled real-time PCR assay for sensitive and specific detection of *Blastocystis* spp. *J. Clin. Microbiol.*
- Stensvold, C.R., Alfellani, M., Clark, C.G., 2012. Levels of genetic diversity vary dramatically between *Blastocystis* subtypes. *Infection. Genetics and Evolution*. <https://doi.org/10.1016/j.meegid.2011.11.002>.
- Stensvold, C.R., Berg, R.P.K.D., Maloney, J.G., Molokin, A., Santin, M., 2023. Molecular characterization of *Blastocystis* and *Entamoeba* of muskoxen and sheep in Greenland. *Int. J. Parasitol.* 53, 673–685. <https://doi.org/10.1016/j.ijpara.2023.05.005>.
- Stensvold, C.R., Clark, C.G., 2020. Pre-empting Pandora's box: *Blastocystis* subtypes revisited. *Trends Parasitol.*
- Stensvold, C.R., Clark, C.G., 2016. Current status of *Blastocystis*: A personal view. *Parasitol. Int.* <https://doi.org/10.1016/j.parint.2016.05.015>.
- Stensvold, C.R., Tan, K.S., Clark, C.G., 2020. *Blastocystis*. *Trends Parasitol.*
- Udonsom, R., Prasertbun, R., Mahittikorn, A., Mori, H., Changbunjong, T., Komalamisra, C., Pintong, A., Sukthana, Y., Popruk, S., 2018. *Blastocystis* infection and subtype distribution in humans, cattle, goats, and pigs in central and western Thailand. *Infection. Genetics and Evolution* 65, 107–111. <https://doi.org/10.1016/j.meegid.2018.07.007>.
- Vega, L., Herrera, G., Muñoz, M., Patarroyo, M.A., Maloney, J.G., Santín, M., Ramírez, J. D., 2021. Gut microbiota profiles in diarrheic patients with co-occurrence of *Clostridioides difficile* and *Blastocystis*. *PLoS One*.
- Vega Romero, L.C., 2020. Cambios en el bacteriana y eucariota intestinal en pacientes con *Blastocystis* y *clostridium difficile*. bachelorThesis. instname:Universidad del Rosario. Universidad del Rosario.
- Villamizar, X., Higuera, A., Herrera, G., Vasquez-A, L.R., Buitron, L., Muñoz, L.M., Gonzalez-C, F.E., Lopez, M.C., Giraldo, J.C., Ramírez, J.D., 2019. Molecular and descriptive epidemiology of intestinal protozoan parasites of children and their pets in Cauca, Colombia: a cross-sectional study. *BMC Infect. Dis.* 19, 190. <https://doi.org/10.1186/s12879-019-3810-0>.
- Wawrzyniak, I., Poirier, P., Viscogliosi, E., Dionigia, M., Texier, C., Delbac, F., Alaoui, H. E., 2013. *Blastocystis*, an unrecognized parasite: an overview of pathogenesis and diagnosis. *Ther. Adv. Infect. Dis.*
- Wylezich, C., Belka, A., Hanke, D., Beer, M., Blome, S., Höper, D., 2019. Metagenomics for broad and improved parasite detection: a proof-of-concept study using swine faecal samples. *Int. J. Parasitol.* 49, 769–777. <https://doi.org/10.1016/j.ijpara.2019.04.007>.
- Xu, Q., Qiao, Q., Gao, Y., Hou, J., Hu, M., Du, Y., Zhao, K., Li, X., 2021. Gut Microbiota and Their Role in Health and Metabolic Disease of Dairy Cow. *Front. Nutr.* 8.
- Yañez, C.M., Hernández, A.M., Sandoval, A.M., Domínguez, M.A.M., Muñoz, S.A.Z., Gómez, J.O.G., 2021. Prevalence of *Blastocystis* and its association with Firmicutes/Bacteroidetes ratio in clinically healthy and metabolically ill subjects. *BMC Microbiol.* 21, 339. <https://doi.org/10.1186/s12866-021-02402-z>.
- Yoshikawa, H., Koyama, Y., Tsuchiya, E., Takami, K., 2016. *Blastocystis* phylogeny among various isolates from humans to insects. *Parasitol. Int.*
- Zhao, G., Hu, X., Liu, T., Hu, R., Yu, Z., Yang, W., Wu, Y., Yu, S., Song, J., 2017. Molecular characterization of *Blastocystis* sp. in captive wild animals in Qinling Mountains. *Parasitol. Res.*